

PEARL

Metabolic basis to Sherpa altitude adaptation

Horscroft, James A.; Kotwica, Aleksandra O.; Laner, Verena; West, James A.; Hennis, Philip J.; Levett, Denny Z.H.; Howard, David J.; Fernandez, Bernadette O.; Burgess, Sarah L.; Ament, Zsuzsanna; Gilbert-Kawai, Edward T.; Vercueil, André; Landis, Blaine D.; Mitchell, Kay; Mythen, Monty G.; Branco, Cristina; Johnson, Randall S.; Feelisch, Martin; Montgomery, Hugh E.; Griffin, Julian L.; Grocott, Michael P.W.; Gnaiger, Erich; Martin, Daniel S.; Murray, Andrew J.

Published in:

Proceedings of the National Academy of Sciences

DOI:

[10.1073/pnas.1700527114](https://doi.org/10.1073/pnas.1700527114)

Publication date:

2017

Link:

[Link to publication in PEARL](#)

Citation for published version (APA):

Horscroft, J. A., Kotwica, A. O., Laner, V., West, J. A., Hennis, P. J., Levett, D. Z. H., Howard, D. J., Fernandez, B. O., Burgess, S. L., Ament, Z., Gilbert-Kawai, E. T., Vercueil, A., Landis, B. D., Mitchell, K., Mythen, M. G., Branco, C., Johnson, R. S., Feelisch, M., Montgomery, H. E., ... Murray, A. J. (2017). Metabolic basis to Sherpa altitude adaptation. *Proceedings of the National Academy of Sciences*, 114(24), 6382-6387.
<https://doi.org/10.1073/pnas.1700527114>

Metabolic basis to Sherpa altitude adaptation

James Horscroft¹, Aleksandra Kotwica¹, Verena Laner², James West¹, Philip Hennis³, Denny Levett³, David Howard³, Bernadette Fernandez⁴, Sarah Burgess¹, Zsuzsanna Ament¹, Edward Gilbert-Kawai³, Andre Vercueil³, Blaine Landis³, Kay Mitchell⁴, Monty Mythen³, Cristina Branco¹, Randall Johnson¹, Martin Feelisch⁴, Hugh Montgomery³, Julian Griffin¹, Michael Grocott⁴, Erich Gnaiger⁵, Daniel Martin³, Andrew J Murray¹

¹University of Cambridge, ²Oroboros Instruments, ³University College London, ⁴University of Southampton, ⁵Innsbruck Medical University

Submitted to Proceedings of the National Academy of Sciences of the United States of America

The Himalayan Sherpas, a human population of Tibetan descent, are highly adapted to life in the hypobaric hypoxia of high altitude. Mechanisms involving enhanced tissue oxygen delivery in comparison with Lowlander populations, have been postulated to play a role in such adaptation. Whether differences in tissue oxygen utilization (i.e. metabolic adaptation) underpin this adaptation is not however known. We sought to address this issue, applying parallel molecular, biochemical, physiological and genetic approaches to the study of Sherpas and native Lowlanders, studied before and during exposure to hypobaric hypoxia on a gradual ascent to Mount Everest Base Camp (5,300 m). When compared with Lowlanders, Sherpas demonstrated a lower capacity for fatty acid oxidation in skeletal muscle biopsies, along with enhanced efficiency of oxygen utilization, improved muscle energetics and protection against oxidative stress. This in part appeared to be related to a putatively advantageous allele for the *PPARA* gene, which was enriched in the Sherpas compared with the Lowlanders. Our findings suggest that metabolic adaptations underpin human evolution to life at high altitude, and could impact upon our understanding of human diseases in which hypoxia is a feature.

metabolism | altitude | skeletal muscle | hypoxia | mitochondria

Introduction

At high altitude, low barometric pressure is accompanied by a fall in the partial pressure of inspired O₂, resulting in *hypobaric hypoxia*. The cellular response to hypoxia is orchestrated by the Hypoxia Inducible Factor (HIF) transcription factors, with HIF-1 α and HIF-2 α respectively mediating responses to short-term and more sustained hypoxia (1). In normoxia, prolyl-hydroxylases target HIF α subunits for destruction (2). Under low O₂ partial pressures, however, HIF-1 α /HIF-2 α are stabilized and dimerize with the nuclear HIF-1 β subunit. This dimer interacts with hypoxia-response elements in promoter regions to increase expression of specific genes, e.g. *EPO* (encoding erythropoietin) and *VEGFA* (vascular endothelial growth factor A) (3).

The Tibetan Plateau has an average altitude of some 4,500 m. Humans were first present on the Plateau ~30,000 years ago, with the earliest permanent settlements appearing 6-9,000 years ago (4) – a period sufficient to drive the natural selection of genetic variants (and associated features) favouring survival and performance in sustained hypoxia (5, 6). Evidence supports the selection of genetic variants encoding components of the hypoxia-inducible factor (HIF) pathway, such as *EPAS1* (encoding HIF-2 α) (7) and *EGLN1* (prolyl-hydroxylase-2, PHD2) (8) in Tibetan populations. One population, the Sherpas, migrated from Tibet to eastern Nepal ~500 years ago and exhibit remarkable physical performance at extreme altitude (9).

Whilst the human adaptive response to hypoxia is incompletely understood, mitigation against the fall in convective O₂ delivery plays an important role. In Lowlanders, increased ventilation and cardiac output, and the production of more O₂-carrying red blood cells help to sustain O₂ delivery and content (10, 11). Likewise, exhaled concentrations of nitric oxide (NO), a key regulator of blood flow, are higher in Tibetans than Lowlanders (12), as are circulating NO metabolites and limb blood flow (13).

The rise in red cell mass in response to hypobaric hypoxia is not as great in Tibetans as in Lowlanders, however (14, 15), suggesting that adaptation involves more than just increased O₂ delivery. In fact, acclimatization also involves alterations in O₂ use. In Lowlander muscle, mitochondrial density declines with sustained exposure to extreme altitude (16-18), whilst exposure to more moderate high altitude is associated with a reprogramming of muscle metabolism (19) even without altered mitochondrial density (20), including downregulation of electron transfer complexes (19) and tricarboxylic acid (TCA) cycle enzymes (21), loss of fatty acid oxidation (FAO) capacity (19, 20) and improved oxidative phosphorylation coupling efficiency (20). Sherpas have lower muscle mitochondrial densities than unacclimatized Lowlanders (22), but little is known of their metabolic adaptation to hypoxia, or any genetic selection which might underpin it. A role has been suggested for peroxisome proliferator-activated receptor alpha (PPAR α), a transcriptional regulator of FAO in liver, heart and muscle. HIF downregulates PPAR α in some tissues (23), whilst there is evidence for selection of variants in its encoding gene (*PPARA*) in some Tibetan subgroups (8, 24). We hypothesized that metabolic adaptation, and PPAR α in particular, play a central role in the Sherpa adaptation to hypobaric hypoxia.

Results and Discussion

Selection of *PPARA* Variants in Sherpas

Lowlander and Sherpa subjects were participants of the research expedition, Xtreme Everest 2 (25). The Lowlanders comprised 10 investigators selected to operate the Everest Base Camp (EBC) laboratory. Sherpas ($n = 15$) were a sex-matched (73%

Significance

A relative fall in tissue oxygen levels (hypoxia) is a common feature of many human diseases including heart failure, lung diseases, anemia and many cancers, and can compromise normal cellular function. Hypoxia also occurs in healthy humans at high altitude due to low barometric pressures. Human populations resident at high altitude in the Himalayas have evolved mechanisms that allow them to survive and perform, including adaptations that preserve oxygen delivery to the tissues. Here we studied one such population, the Sherpas, and found metabolic adaptations, underpinned by genetic differences, which allow their tissues to use oxygen more efficiently, thereby conserving muscle energy levels at high altitude, and possibly contributing to the superior performance of elite climbing Sherpas at extreme altitudes.

Reserved for Publication Footnotes

137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204




Fig. 1. Subject genetics, ascent profile, arterial blood O₂ saturation, muscle hypoxia and circulating NO metabolites. A) Genotypes of Lowlanders and Sherpas at 3 *PPARA* SNPs - subjects homozygous for the putatively advantageous allele in black, heterozygous subjects in gray and subjects homozygous for the non-advantageous allele in white (digits in segments refer to number of subjects with genotype); B) Ascent profile including timing of biopsies; C) Arterial hemoglobin-O₂ saturations; D) Muscle *VEGFA* expression, and E-H) plasma nitrogen oxides in Lowlanders (L) and Sherpas (S) at baseline (B) and early (A1) and late (A2) altitude. Mean \pm SEM ($n = 4-15$). $\dagger P \leq 0.05$; $\dagger\dagger P \leq 0.001$ B vs A1 within cohort. $\Delta P \leq 0.05$ A1 vs A2 within cohort.

male, *cf.* 70% in Lowlanders) and age-matched (26.8 ± 1.2 yr, *cf.* 28.0 ± 1.6 yr in Lowlanders) group living in Kathmandu and the




Fig. 2. Fatty acid oxidation and regulation in muscle. A) *PPARA* expression; B) *CPT1B* expression; C) HADH activity; D) Oxidative phosphorylation with octanoylcarnitine:malate (FAOP); E) Total carnitine; F) Long chain/total carnitine ratio in Lowlanders and Sherpas. Gene expression and carnitine levels are expressed relative to Lowlanders at baseline. Mean \pm SEM ($n = 6-13$). $*P \leq 0.05$; $**P \leq 0.01$ Lowlanders vs Sherpas at baseline. $\dagger P \leq 0.05$ baseline vs altitude within cohort.

Solkhumbu and Rolwaling valleys. No subject ascended higher than 4,200 m in the 3 months preceding the trek, nor above 2,500 m in the preceding 3 weeks. In addition, Sherpas presented evidence of sole Sherpa ancestry for 2 generations (i.e. 4 Sherpa grandparents). The frequency of putatively advantageous *PPARA* alleles (8) was higher in Sherpas than Lowlanders (Fig. 1A; Table S1), with genotype frequencies of the cohorts being significantly different at 2 single nucleotide polymorphisms (SNPs), rs6520015 and rs7292407 ($P = 0.0091$), though not rs9627403. This reflected patterns reported in some other Tibetan groups (26).

Muscle Hypoxia and Circulating NO Metabolites

Baseline testing, including blood sampling, muscle biopsy sampling, high-resolution respirometry of permeabilized muscle fibers and oral glucose tolerance tests (OGTT) took place in London (35 m) for Lowlanders and Kathmandu (1,300 m) for Sherpas (25). All subjects then followed an identical ascent (Fig. 1B) from Kathmandu to EBC (5,300 m) whereupon further testing took place at an early timepoint (A1; 15-20 d post-departure for Lowlanders, 11-12 d for Sherpas), and a late timepoint (A2; 54-59 d post-departure) for Lowlanders only. At the time of sampling, both groups had passed through the acute phase of hypoxic exposure (<24 h) (1) and had been sufficiently exposed to chronic hypoxia for acclimatization to have occurred. Indeed, arterial hemoglobin-O₂ saturations were similarly low in both groups (Fig. 1C), whilst muscle expression of the HIF-target *VEGFA* increased in all subjects (Fig. 1D), indicating a molecular response to hypoxia. Following measurements at A1, the Lowlanders remained at EBC for 2 months to carry out research, presenting an opportunity to collect data pertaining to longer-term metabolic acclimatization. Interestingly, *VEGFA*

205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272

273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340




Fig. 3. TCA intermediates and activity in muscle. A) Citrate synthase activity and B-I) TCA cycle intermediates in Lowlanders and Sherpas. Metabolite levels are expressed relative to Lowlanders at baseline. Mean \pm SEM ($n = 7-14$). * $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$ Lowlanders vs Sherpas at baseline. † $P \leq 0.05$; †† $P \leq 0.01$; baseline vs altitude within cohort.

expression was no longer elevated by this timepoint, suggesting further acclimatization had occurred.





Fig. 4. Mitochondrial oxygen consumption, efficiency and uncoupling protein expression. A) N-OXPHOS (GM_P), B) S-ETS capacity (S_E) and C) NS-OXPHOS capacity (GMS_P) in permeabilized muscle fibers from Lowlanders and Sherpas. D) Octanoylcarnitine&malate-supported LEAK (FAO_L) and E) OXPHOS coupling efficiency. F) Muscle *UCP3* expression relative to Lowlanders at baseline. Mean \pm SEM ($n = 7-11$). ** $P \leq 0.01$; *** $P \leq 0.001$ Lowlander vs Sherpas at baseline. † $P \leq 0.05$; †† $P \leq 0.01$ baseline vs altitude within cohort. $\Delta P \leq 0.05$; $\Delta\Delta P \leq 0.01$ altitude 1 vs 2 within cohort.

To our surprise, there were no differences in circulating N-nitrosamine (RNNO), S-nitrosothiol (RSNO), nitrate (NO_3^-) or nitrite (NO_2^-) concentrations between Lowlanders and Sherpas at baseline (Fig. 1E-H). In Lowlanders, a transient increase in plasma RNNO levels occurred upon arrival at EBC ($P < 0.05$) but disappeared by the later timepoint (Fig. 1E). In Sherpas, plasma nitrate levels fell at altitude ($P < 0.05$; Fig. 1G) and nitrite levels increased ($P < 0.05$; Fig. 1H), whilst in Lowlanders nitrite levels fell by the later timepoint ($P < 0.05$). The absence of large differences in NO metabolites between the groups at baseline or at altitude, suggested an adaptive phenotype in Sherpas that is distinct from other Tibetan highlanders (13).

Lower Fatty Acid Oxidation Capacity in Sherpas

Skeletal muscle biopsies revealed marked differences in gene expression and FAO capacity between Sherpas and Lowlanders. Expression of *PPARA* mRNA was 48% lower in Sherpas than Lowlanders ($P < 0.05$; Fig. 2A), thus the putatively advantageous *PPARA* allele is associated with diminished expression. Correspondingly, expression of the *PPARA* target *CPT1B* was 32% lower in Sherpas at baseline compared with Lowlanders ($P < 0.05$; Fig. 2B). The *PPARA* gene contains 139 SNPs. rs6520015 is one of the tagging SNPs reported by Simonson *et al* (8), however it appears to be a non-coding variant. It is thus uncertain whether the SNP itself affects transcriptional regulation, or whether it tags a functional variant elsewhere, modifying expression or mRNA stability. Ascent to EBC did not alter *PPARA* expression in either group, yet despite this *CPT1B* expression decreased by 44% in



341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408



Lowlanders ($P < 0.05$) but did not decrease further in Sherpas. This suggests that the Lowlander response to hypoxia involves decreased PPAR α transcriptional activity without changes in PPARA expression, similar to hypoxic rat skeletal muscle (27).

Gene expression changes do not necessarily reflect protein levels or activity, therefore we measured activity of the β -oxidation enzyme 3-hydroxyacyl-CoA dehydrogenase (HADH), finding it to be 27% lower in Sherpas than Lowlanders at baseline ($P < 0.05$), and not changing in either group following ascent (Fig. 2C). Moreover, fatty acid oxidative phosphorylation capacity (FAO $_P$) was measured as the oxygen flux in saponin-permeabilized muscle fibers with octanoylcarnitine, malate and ADP, using high-resolution respirometry (28). FAO $_P$ was 24% lower in Sherpas than Lowlanders at baseline ($P < 0.01$), and did not change in either group following ascent (Fig. 2D, Fig. S1). *Ex vivo* measurements may be particular to assay conditions used, therefore we also measured muscle metabolite levels to indicate changes in metabolism *in vivo*.

Total carnitine concentrations decreased in Lowlanders with time spent at altitude ($P < 0.05$), though were not significantly different to those in Sherpas at baseline (Fig. 2E). The ratio of long chain acylcarnitines to total carnitines, however, increased in Lowlanders with time at altitude ($P < 0.05$; Fig. 2F), suggesting incomplete FAO results in accumu-



lation of potentially-harmful lipid intermediates (29). In Sherpa muscle, however, the long chain acylcarnitine to total carnitine ratio was lower than in Lowlanders at baseline ($P < 0.05$), perhaps resulting from lower expression of CPT-1. In further contrast with Lowlanders, the long chain acylcarnitine to total carnitine ratio remained low in Sherpa muscle at altitude.

TCA Cycle Regulation at High Altitude

We therefore sought to understand whether there were differences between the populations in other aspects of mitochondrial metabolism. The TCA cycle enzyme citrate synthase (CS) is a candidate marker of mitochondrial content in human muscle (30). At baseline, Sherpas had a 26% lower muscle CS activity than Lowlanders ($P < 0.05$; Fig. 3A), in agreement with findings of 17-33% lower mitochondrial volume density in Sherpa *vastus lateralis* compared with Lowlanders (22). In accordance with lower CS activity, concentrations of 6- and 5-carbon intermediates downstream of CS (citrate, aconitate, isocitrate, α -ketoglutarate) were lower in Sherpas than Lowlanders ($P < 0.001$). However, concentrations of 4-carbon intermediates (succinate, fumarate, malate, oxaloacetate) were not different (Fig 3B-I). This suggests an alternative strategy to supply the TCA cycle with succinate. Intriguingly, recent analysis of a large SNP dataset from low and high altitude-adapted populations in the Americas and Asia (31) aimed to identify pathways of convergent evolution, and highlighted fatty acid ω -oxidation as the most significant cluster of overlapping gene sets between high altitude groups (32). ω -oxidation, is normally a minor pathway in vertebrates, becoming more important when β -oxidation is defective (33), and through successive cycles oxidizes fatty acids to adipate and succinate in the endoplasmic reticulum, after which succinate enters the mitochondria with anaplerotic regulation of the TCA cycle (34).

Upon ascent to altitude, 6- and 5-carbon TCA cycle intermediates increased in Sherpa muscle ($P < 0.05$; Fig. 3B-E), suggesting improved coupling of intermediary metabolism, TCA cycle and oxidative phosphorylation. In Lowlanders, however, citrate, aconitate and isocitrate decreased at altitude ($P < 0.05$; Fig. 3B-D), despite no significant change in CS activity, perhaps reflecting impairments upstream. Interestingly, α -ketoglutarate concentrations were maintained in Lowlanders at altitude (Fig. 3E), despite decreased succinate downstream, which could be explained by the fall in both α -ketoglutarate dehydrogenase and isocitrate

dehydrogenase, reported previously in Lowlanders following an identical ascent to EBC (21). α -ketoglutarate plays regulatory roles in hypoxia, including a suppression of HIF stabilization (35), but also supporting glutathione synthesis (36). Taken together, these results indicate different TCA cycle regulation in Sherpas and Lowlanders. The replete TCA cycle of Sherpas at altitude contrasts sharply with the depletion of TCA cycle intermediates in Lowlanders, and suggests a coupling of the TCA cycle in Sherpa muscle to their distinct intermediary substrate metabolism.

Greater Mitochondrial Coupling Efficiency in Sherpas

To further understand whether mitochondrial function differs between Sherpas and Lowlanders, we used high-resolution respirometry, to probe electron transfer system (ETS) capacity and coupling efficiency in permeabilized muscle fibers. At baseline, there was no significant difference between the two groups in OXPHOS or ETS capacities with either malate and glutamate (N-pathway through Complex I) or succinate as substrates (S-pathway through Complex II; Fig. 4A,B; Fig. S2), but Sherpas had a lower OXPHOS capacity with malate, glutamate and succinate combined to reconstitute TCA cycle function (NS-pathway; $P < 0.01$; Fig. 4C). There were no early changes in either group upon ascent. By the later timepoint however, succinate-linked respiration had fallen in Lowlanders ($P < 0.05$), consistent with previous findings of decreased succinate dehydrogenase (Complex II) levels in subjects with sustained exposure $>5,300$ m (21).

In addition, we measured muscle fiber respiration in the absence of ADP (LEAK), i.e. O_2 consumption without ADP phosphorylation. Expressing LEAK relative to OXPHOS capacity, it is possible to calculate OXPHOS coupling efficiency (37, 38). At baseline, Sherpa muscle mitochondria had lower LEAK respiration and greater coupling efficiency than Lowlander mitochondria ($P < 0.001$; Fig. 4D,E), indicating more efficient use of O_2 . Upon ascent to EBC and with sustained time at altitude, LEAK decreased in Lowlanders ($P < 0.01$), though it remained higher than in Sherpas (Fig. 4D), and coupling efficiency improved ($P < 0.05$; Fig. 4E). In Sherpas at altitude, LEAK did not change although coupling efficiency decreased ($P < 0.01$). One possible explanation for these differences in coupling efficiency might be the altered expression of uncoupling protein 3 (UCP3). UCP3 is a transcriptional target of PPAR α and lower UCP3 levels at altitude might improve the efficiency of O_2 utilization. In previous studies, however, muscle UCP3 expression increased with acute hypoxia (17, 39), which may offer some protective benefit considering its possible role as an antioxidant (39). Notably though, UCP3 levels decreased with more sustained exposure to extreme altitude (17). Here, UCP3 was upregulated in Sherpas at altitude in association with decreased coupling efficiency ($P < 0.05$; Fig. 4F). However, UCP3 expression also increased in Lowlanders in the short-term ($P < 0.01$) in whom there was decreased LEAK respiration. Moreover, UCP3 expression returned to baseline in Lowlanders with longer-term exposure with no further change in LEAK respiration. Overall, our results indicate that Sherpa muscle mitochondria are characterized by a lower OXPHOS capacity and greater, albeit declining, efficiency, whilst in Lowlanders OXPHOS efficiency improved with acclimatization.

Glycolysis and Glucose Metabolism

Next we investigated the capacity to derive cellular energy via glycolysis, which is increased in hypoxic cells (40), as this may allow ATP levels to be maintained when O_2 is limited. Hexokinase activity was the same in both groups at baseline, and did not change at altitude (Fig. 5A), however lactate dehydrogenase (LDH) activity was 48% higher in Sherpa muscle than in Lowlanders ($P < 0.05$), indicating greater capacity for anaerobic lactate production (Fig. 5B). Fasting blood glucose was the same in Sherpas and Lowlanders at baseline, and decreased upon ascent in Lowlanders ($P < 0.01$; Fig. 5C), who also showed faster clearance of glucose during an OGTT ($P < 0.001$; Fig. 5D) in agree-

ment with previous reports (41). In Sherpas, however, there was no indication of altered glucose homeostasis. Meanwhile, over time at altitude glycolytic intermediates increased in Lowlander muscle (Fig. 5E) with increased glucose-6-phosphate/fructose-6-phosphate and 2-phosphoglycerate/3-phosphoglycerate (Table S2). In contrast, total glycolytic intermediates did not change in Sherpa muscle, although 2-phosphoglycerate/3-phosphoglycerate decreased. These findings, might to some extent be explained by altered HIF activities. Many genes encoding glycolytic enzymes are upregulated by HIF-1 (42), whilst hypoglycemia is seen in Chuvash polycythemia, an autosomal recessive disorder in which HIF degradation is impaired (43). Taken together, our findings suggest an increased reliance on glucose by Lowlanders under resting conditions at altitude compared with Sherpas, but a greater capacity for lactate production in Sherpas which may prove effective upon exertion.

Energetics and Oxidative Stress

Finally, to understand the implications of Sherpa metabolic adaptation we investigated muscle energetics and redox homeostasis. Lowlanders at altitude showed progressive loss of muscle phosphocreatine (PCr; $P < 0.001$; Fig. 6A), indicating a loss of energetic reserve, which may relate to downregulation of muscle creatine kinase, as reported previously (21). By contrast, in Sherpa muscle, PCr increased at altitude ($P < 0.01$). Similarly, Sherpa muscle ATP levels, which were lower than in Lowlanders at baseline ($P < 0.05$), increased at altitude ($P < 0.001$; Fig. 6B), illustrating that Sherpa metabolism is better suited to maintaining muscle energetics at altitude than Lowlander metabolism in either the short-term or following acclimatization. Moreover, with short-term exposure, markers of oxidative stress (reduced/oxidized glutathione and methionine sulfoxide) increased in Lowlander muscle, but not Sherpa muscle (Fig. 6C,D), indicating superior redox homeostasis in the Sherpas. Antioxidant protection may represent another outcome of convergent evolution, having been reported in Andean subjects in association with protection of fetal growth (44), whilst glutathione levels are raised in Chuvash polycythemia suggesting a possible role for HIF activation (45).

Conclusions

It has long been suspected that Sherpa people are better adapted to life at high altitude than Lowlanders (46). Recent findings have suggested a genetic basis to adaptation in populations around the world (6), and here we show that Sherpas have a metabolic adaptation associated with improved muscle energetics and protection against oxidative stress. Genetic selection on the PPARA gene is associated with decreased expression, and thus lower fatty acid β -oxidation and improved mitochondrial coupling compared with Lowlanders, with a possible compensatory increase in fatty acid ω -oxidation. Sherpas also have a greater capacity for lactate production. With acclimatization to altitude, Lowlanders accumulate potentially-harmful lipid intermediates in muscle as a result of incomplete β -oxidation, alongside depletion of TCA cycle intermediates, accumulation of glycolytic intermediates, a loss of PCr despite improved mitochondrial coupling, and a transient increase in oxidative stress markers. In Sherpas, however, there are remarkably few changes in intermediary metabolism at altitude, but increased TCA cycle intermediates and PCr and ATP levels, with no sign of oxidative stress.

Genetic selection, by definition, requires an increased likelihood of advantageous gene variants being passed on to offspring. This might occur if the disadvantageous variant is associated with poorer survival to reproductive age and beyond, including greater fetal/neonatal mortality. Evidence supports precisely such effects with fetal growth at altitude being poorer in Lowlander populations than many native highlanders (47), including Tibetans (48) and Sherpas (49). Likewise, gene variants may affect

681 survival through childhood or fecundity/fertility in the hypoxic
682 environment. We cannot speculate on the mechanism by which
683 *PPARA* variants prove advantageous, however PPAR isoforms are
684 expressed in the placenta (50) and influence female reproductive
685 function (51). It would be of interest to seek association of the
686 *PPARA* variants with birth weight and measures of placentation
687 in high altitude natives and Lowlanders exposed to hypoxia.

688 Our findings suggest a metabolic basis to Sherpa adaptation,
689 which may permit the population to survive and perform at
690 high altitude. Such adaptations may also underpin the superior
691 performance of elite climbing Sherpas at extreme high altitude.

693 Materials and Methods

694 Subjects were selected from the participants of Xtreme Everest 2 (25).
695 All Lowlanders were born and lived below 1,000 m, not descended from
696 a high altitude-dwelling population and of European (Caucasian) origin.
697 Subjects gave written consent, and underwent medical screening. All proto-
698 cols were approved by UCL Research Ethics Committee and Nepal Health
699 Research Council. Vastus lateralis biopsies were taken from the mid-thigh,

1. Koh MY, Powis G (2012) Passing the baton: the HIF switch. *Trends Biochem Sci* 37(9):364-372.
2. Willam C, Nicholls LG, Ratcliffe PJ, Pugh CW, Maxwell PH (2004) The prolyl hydroxylase enzymes that act as oxygen sensors regulating destruction of hypoxia-inducible factor alpha. *Adv Enzyme Regul* 44:75-92.
3. Semenza GL (2012) Hypoxia-inducible factors in physiology and medicine. *Cell* 148(3):399-408.
4. Aldenderfer M (2011) Peopling the Tibetan plateau: insights from archaeology. *High Alt Med Biol* 12(2):141-147.
5. Beall CM (2007) Two routes to functional adaptation: Tibetan and Andean high-altitude natives. *Proc Natl Acad Sci USA* 104 Suppl 1:8655-8660.
6. Bigham AW, Lee FS (2014) Human high-altitude adaptation: forward genetics meets the HIF pathway. *Genes Dev* 28(20):2189-2204.
7. Beall CM, et al. (2010) Natural selection on EPAS1 (HIF2alpha) associated with low hemoglobin concentration in Tibetan highlanders. *Proc Natl Acad Sci USA* 107(25):11459-11464.
8. Simonson TS, et al. (2010) Genetic evidence for high-altitude adaptation in Tibet. *Science* 329(5987):72-75.
9. Gilbert-Kawai ET, Milledge JS, Grocott MP, Martin DS (2014) King of the mountains: Tibetan and Sherpa physiological adaptations for life at high altitude. *Physiology* 29(6):388-402.
10. Peacock AJ (1998) ABC of oxygen: oxygen at high altitude. *BMJ* 317(7165):1063-1066.
11. Grocott MP, et al. (2009) Arterial blood gases and oxygen content in climbers on Mount Everest. *N Engl J Med* 360(2):140-149.
12. Beall CM, et al. (2001) Pulmonary nitric oxide in mountain dwellers. *Nature* 414(6862):411-412.
13. Erzurum SC, et al. (2007) Higher blood flow and circulating NO products offset high-altitude hypoxia among Tibetans. *Proc Natl Acad Sci USA* 104(45):17593-17598.
14. Winslow RM, et al. (1989) Different hematologic responses to hypoxia in Sherpas and Quechua Indians. *J Appl Physiol* 66(4):1561-1569.
15. Beall CM, et al. (1998) Hemoglobin concentration of high-altitude Tibetans and Bolivian Aymara. *Am J Phys Anthropol* 106(3):385-400.
16. Hoppeler H, Howald H, & Cerretelli P (1990) Human muscle structure after exposure to extreme altitude. *Experientia* 46(11-12):1185-1187.
17. Levett DZ, et al. (2012) Acclimatization of skeletal muscle mitochondria to high-altitude hypoxia during an ascent of Everest. *FASEB J* 26(4):1431-1441.
18. Murray AJ, Horscroft JA (2016) Mitochondrial function at extreme high altitude. *J Physiol* 594(5):1137-1149.
19. Horscroft JA, Murray AJ (2014) Skeletal muscle energy metabolism in environmental hypoxia: climbing towards consensus. *Extrem Physiol Med* 3(1):19.
20. Jacobs RA, et al. (2012) Twenty-eight days at 3454-m altitude diminishes respiratory capacity but enhances efficiency in human skeletal muscle mitochondria. *FASEB J* 26(12):5192-5200.
21. Levett DZ, et al. (2015) Changes in muscle proteomics in the course of the Caudwell Research Expedition to Mt. Everest. *Proteomics* 15(1):160-171.
22. Kayser B, Hoppeler H, Claassen H, Cerretelli P (1991) Muscle structure and performance capacity of Himalayan Sherpas. *J Appl Physiol* 70(5):1938-1942.
23. Narravula S, Colgan SP (2001) Hypoxia-inducible factor 1-mediated inhibition of peroxisome proliferator-activated receptor alpha expression during hypoxia. *J Immunol* 166(12):7543-7548.
24. Peng Y, et al. (2011) Genetic variations in Tibetan populations and high-altitude adaptation at the Himalayas. *Mol Biol Evol* 28(2):1075-1081.
25. Gilbert-Kawai E, et al. (2015) Design and conduct of Xtreme Everest 2: An observational cohort study of Sherpa and lowlander responses to graduated hypobaric hypoxia. *F1000Res* 4:90.
26. Ge RL, et al. (2012) Metabolic insight into mechanisms of high-altitude adaptation in Tibetans. *Mol Genet Metab* 106(2):244-247.
27. Horscroft JA, Burgess SL, Hu Y, Murray AJ (2015) Altered Oxygen Utilisation in Rat Left Ventricle and Soleus after 14 Days, but Not 2 Days, of Environmental Hypoxia. *PLoS one* 10(9):e0138564.
28. Pesta D, Gnaiger E (2012) High-resolution respirometry: OXPHOS protocols for human cells

700 muscle fibers prepared for respirometry (28) and respiration measured using
701 substrate-uncoupler-inhibitor titrations (Tables S3, S4). Enzyme activities
702 were assayed as described (27). RNA was extracted and Taqman® assays
703 used to analyse gene expression (Table S5). For metabolite analysis, a
704 methanol/chloroform extraction (52) was followed by liquid chromatogra-
705 phy mass spectrometry (LC-MS). OGTTs were carried out on fasted subjects
706 on the day after biopsies. Blood plasma NO metabolites were quantified as
707 described (53). Genomic DNA was isolated from whole blood and *PPARA* SNPs
708 genotyped using TaqMan® for allelic discrimination (Applied Biosystems, UK;
709 Table S1). To compare cohorts at baseline, an unpaired two-tailed Student's t-
710 test was used (significance at P ≤ 0.05). Genotype frequencies were compared
711 using a Chi-squared test. To assess the effects of altitude, a one-way ANOVA
712 with repeated measures was used. Post-hoc pairwise comparisons were
713 carried out with a Tukey correction.

714 Acknowledgements

715 The work was supported by PhD studentships from the BBSRC to JH
716 (BB/F016581/1) and British Heart Foundation to AK (FS/09/050), an Academic
717 Fellowship to AM from the Research Councils UK (EP/E500552/1), a Physio-
718 logical Society grant and support from Oroboros Instruments. JG thanks the
719 MRC (MC UP A90 1006) and AB Sciex. MF thanks the MRC and Faculty of
720 Medicine, Southampton University. For full acknowledgements see SI.

- and permeabilized fibers from small biopsies of human muscle. *Methods Mol Biol* 810:25-58.
29. Koves TR, et al. (2008) Mitochondrial overload and incomplete fatty acid oxidation contribute to skeletal muscle insulin resistance. *Cell Metab* 7(1):45-56.
30. Kayser B, et al. (1996) Muscle ultrastructure and biochemistry of lowland Tibetans. *J Appl Physiol* 81(1):419-425.
31. Bigham A, et al. (2010) Identifying signatures of natural selection in Tibetan and Andean populations using dense genome scan data. *PLoS Genet* 6(9):e1001116.
32. Foll M, Gaggiotti OE, Daub JT, Vatsiou A, Excoffier L (2014) Widespread signals of convergent adaptation to high altitude in Asia and America. *Am J Hum Genet* 95(4):394-407.
33. Wanders RJ, Komen J, Kemp S (2011) Fatty acid omega-oxidation as a rescue pathway for fatty acid oxidation disorders in humans. *FEBS J* 278(2):182-194.
34. Nelson DL, Cox MM, Lehninger AL (2008) *Principles of Biochemistry* (W.H. Freeman, New York)
35. MacKenzie ED, et al. (2007) Cell-permeating alpha-ketoglutarate derivatives alleviate pseudohypoxia in succinate dehydrogenase-deficient cells. *Mol Cell Biol* 27(9):3282-3289.
36. Chinopoulos C (2013) Which way does the citric acid cycle turn during hypoxia? The critical role of alpha-ketoglutarate dehydrogenase complex. *J Neurosci Res* 91(8):1030-1043.
37. Gnaiger E, et al. (2015) Mitochondrial coupling and capacity of oxidative phosphorylation in skeletal muscle of Inuit and Caucasians in the arctic winter. *Scand J Med Sci Sports* 25 Suppl 4:126-134.
38. Gnaiger E (2014) Mitochondrial pathways and respiratory control. An introduction to OXPHOS analysis. *Mitochondr Physiol Network* 19(12).
39. Anedda A, et al. (2013) The transcription factor Nrf2 promotes survival by enhancing the expression of uncoupling protein 3 under conditions of oxidative stress. *Free Radic Biol Med* 61:395-407.
40. Murray AJ (2009) Metabolic adaptation of skeletal muscle to high altitude hypoxia: how new technologies could resolve the controversies. *Genome Med* 1(12):117.
41. Woolcott OO, Ader M, Bergman RN (2015) Glucose homeostasis during short-term and prolonged exposure to high altitudes. *Endocr Rev* 36(2):149-173.
42. Semenza GL, Roth PH, Fang HM, Wang GL (1994) Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1. *J Biol Chem* 269(38):23757-23763.
43. McClain DA, et al. (2013) Decreased serum glucose and glycosylated hemoglobin levels in patients with Chuvash polycythemia: a role for HIF in glucose metabolism. *J Mol Med* 91(1):59-67.
44. Julian CG, et al. (2012) Potential role for elevated maternal enzymatic antioxidant status in Andean protection against altitude-associated SGA. *J Matern Fetal Neonatal Med* 25(8):1233-1240.
45. Sergueeva AI, et al. (2008) Elevated homocysteine, glutathione and cysteinylglycine concentrations in patients homozygous for the Chuvash polycythemia VHL mutation. *Haematologica* 93(2):279-282.
46. Lahiri S, Milledge JS (1965) Sherpa physiology. *Nature* 207(997):610-612.
47. Moore LG, Charles SM, Julian CG (2011) Humans at high altitude: hypoxia and fetal growth. *Respir Physiol Neurobiol* 178(1):181-190.
48. Moore LG, Young D, McCullough RE, Droma T, Zamudio S (2001) Tibetan protection from intrauterine growth restriction (IUGR) and reproductive loss at high altitude. *Am J Hum Biol* 13(5):635-644.
49. Smith C (1997) The effect of maternal nutritional variables on birthweight outcomes of infants born to Sherpa women at low and high altitudes in Nepal. *Am J Hum Biol* 9(6):751-763.
50. Jawerbaum A, Capobianco E (2011) Review: Effects of PPAR activation in the placenta and the fetus: implications in maternal diabetes. *Placenta* 32 Suppl 2:S212-217.
51. Bogacka I, Kurzynska A, Bogacki M, Chojnowska K (2015) Peroxisome proliferator-activated receptors in the regulation of female reproductive functions. *Folia Histochem Cytobiol* 53(3):189-200.
52. Roberts LD, et al. (2011) The contrasting roles of PPARdelta and PPARgamma in regulating the metabolic switch between oxidation and storage of fats in white adipose tissue. *Genome Biol* 12(8):R75.
53. Rassaf T, Bryan NS, Kelm M, Feelisch M (2002) Concomitant presence of N-nitroso and S-nitroso proteins in human plasma. *Free Radic Biol Med* 33(11):1590-1596.