



PEARL

Impacts of polyethylene microplastics on the microalga, *Spirulina* (*Arthrospira platensis*)

Abbasi, Sajjad; Amiranipour, Sahar; Karimi, Javad; Mohsenzadeh, Sasan; Turner, Andrew

Published in:
Environmental Pollution

DOI:
[10.1016/j.envpol.2023.121611](https://doi.org/10.1016/j.envpol.2023.121611)

Publication date:
2023

Link:
[Link to publication in PEARL](#)

Citation for published version (APA):

Abbasi, S., Amiranipour, S., Karimi, J., Mohsenzadeh, S., & Turner, A. (2023). Impacts of polyethylene microplastics on the microalga, *Spirulina* (*Arthrospira platensis*). *Environmental Pollution*, 327(0). <https://doi.org/10.1016/j.envpol.2023.121611>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 **Impacts of polyethylene microplastics on the microalga, Spirulina**
2 **(*Arthrospira platensis*)**

3

4 Sajjad Abbasi ^{1,3}, Sahar Amiranipour², Javad Karimi*^{2,3}, Sasan Mohsenzadeh ², Andrew
5 Turner ⁴

6 ¹Department of Earth Sciences, School of Science, Shiraz University, Shiraz 71454, Iran

7 ²Department of Biology, School of Science, Shiraz University, Shiraz, 71454, Iran

8 ³Centre for Environmental Studies and Emerging Pollutants (ZISTANO), Shiraz University,
9 Shiraz, 714545, Iran

10 ⁴School of Geography, Earth and Environmental Sciences, University of Plymouth, PL4 8AA,
11 UK


12

13 <https://doi.org/10.1016/j.envpol.2023.121611>

14

15 **Accepted 8 April 2023**

16 **Graphical Abstract**



17

18

19 **Abstract**

20 Microalgae play a critical role in the food web and biogeochemical cycling and produce compounds
21 that are commercially exploited. However, their reactions and responses to microplastic
22 contamination are not well understood. In this study, the widely distributed and commercially
23 important cyanobacterium, *Spirulina (Arthrospira platensis)*, was exposed to different
24 concentrations (1 to 100 mg L⁻¹) of low-density polyethylene microplastics (< 5 μm) over a 20-d
25 period. Various end-points were combined with different microscopic techniques in order to
26 examine physiological and biochemical effects and interactions between the plastic and microalga.
27 Growth rate and photosynthetic activity decreased with increasing microplastic concentration, and a
28 maximum inhibition ratio of about 9% was calculated from optical density measurements. Plastic
29 concentrations above 10 mg L⁻¹ resulted in oxidative stress and the intracellular production of
30 proline. Fragmentation and swelling of trichomes and attachment of microplastics was observed in
31 the exposures, and microplastics appeared to adhere or aggregate around fragmented or
32 fragmenting regions. The latter effect may indicate trichome weakening by microplastics or their
33 concentration around cytosolic debris; nevertheless, it provides a potential mechanism for
34 internalisation of small particles. Although unrealistically high concentrations of well-defined

35 microplastics have been employed, relatively small disruptions at the population level incurred by
36 lower concentrations could have more serious implications for ecosystem services and functioning.

37

38 **Keywords:** cyanobacteria; growth inhibition; photosynthesis; nanoplastics; fragmentation;
39 ecosystem services

40

41 1. Introduction

42 Microalgae are small, photosynthetic, autotrophic organisms that exist individually or in chains or
43 groups and include eukaryotes, diatoms, dinoflagellates and prokaryotic cyanobacteria. Despite
44 ranging in size from only a few μm to several hundred μm , their abundance, efficient biological
45 fixation of carbon and rapid growth rates mean that microalgae consume significant quantities of
46 CO_2 and produce at least one-half of the planet's atmospheric oxygen (Gigova and Marinova, 2016).
47 Coupled with their role in nutrient recycling, microalgae are also a critical link in the food chain.

48 Microalgae also produce a variety of secondary metabolites, such as pigments, dyes, antioxidants
49 and polysaccharides, with structures and activities generally not encountered in other organisms.
50 Consequently, microalgae have attracted commercial and industrial interest in various sectors that
51 include animal husbandry, pharmaceuticals, cosmetics and food production (Priyadarshani and Rath,
52 2012; Skjånes et al., 2017). Among the most widely used microalgae in this respect are
53 *Cyanophyceae* (blue-green algae), *Chlorophyceae*, *Bacillariophyceae* and *Chrysophyceae* (Mobin and
54 Alam, 2017).

55 Microalgae are often exposed to biotic and abiotic stressors and it is critical, therefore, to
56 understand their reactions and responses. One type of aquatic contaminant that has gained
57 considerable interest recently is microplastics, or primary and secondary plastics below 5 mm in size.
58 However, a review by Prata et al. (2019) concluded that the effects of microplastics on microalgae
59 have seldom been determined and that experimental results provide no consensus. For example, the
60 same polymer can promote, inhibit or have no effect on growth, depending on the precise
61 experimental conditions and species employed.

62 Spirulina are filamentous, blue-green cyanobacteria that are found widely in soil, marshes,
63 freshwater, seawater and thermal springs, but thrive in warm, saline, alkaline environments with
64 high levels of insolation. *Arthrospira platensis* is one of the most common and important species in
65 the genera (*Spirulina* and *Arthrospira*), and being easy to harvest and process and with a high
66 nutrient content is popular in the human health food industry and as an aquaculture feed additive
67 (Habib et al., 2008). Despite extensive documentation in scientific research and in public health and
68 food security literature, however, its physiological or biochemical response to microplastic
69 contamination is not well understood. Specifically, exposure to microplastics of different
70 composition appears to inhibit the growth of *Spirulina* sp. and alter biochemical composition and
71 promote the production of extracellular polymeric substances (Abed et al., 2021; Hadiyanto et al.,
72 2021; 2022).

73 In the present study, we investigate the effects of different concentrations of microplastics
74 constructed of polyethylene, one of the most widely used polymers, on the growth, photosynthesis
75 and production of reactive oxygen species (ROS) of *A. platensis*. We combine established
76 methodologies and end-points with Raman spectroscopy and microscopic imagery in order to
77 explore possible mechanisms, morphological changes and plastic-organism interactions involved in
78 these effects and with microalgae more generally.

79

80

81 **2. Methods**

82 *2.1. Materials and exposure conditions*

83 *Spirulina* microalgae, *Spirulina (Arthrospira) platensis*, were cultured in the Department of Biology at
84 Shiraz University. All reagents were purchased from Merck or Sigma-Aldrich and irregularly-shaped
85 polyethylene (PE) microplastics, derived from milling and sieving pure low-density polyethylene
86 (LDPE), were sourced from Torun University, Poland. Scanning electron microscopy (see below)
87 revealed maximum and median particle diameters of about 8 μm and 2.5 μm , respectively, meaning
88 that particles are at the lower end of the size spectrum of microplastics (conventionally defined as 1
89 μm to 5 mm). Microalgae were maintained at 30 ± 2 °C under fluorescent lighting 3500 lux and with
90 a cultivation cycle of 20 d in Zarrouks medium, prepared from ACS or analytical grade salts dissolved
91 in distilled, sterilised water and whose pH was 9.5 and salinity was about 27.

92 Twenty-day exposures of ~ 75 mg *A. platensis* in 1 L Zarrouks medium were performed under the
93 conditions above in a series of 1-L PET cylinders aerated and agitated with individual air stones and
94 stirrers. Exposures ($n = 16$) consisted of quadruplicates of a control (no microplastics), and
95 microplastic concentration spanning two orders of magnitude on a mass basis (1 mg L⁻¹ PE, 10 mg L⁻¹
96 PE and 100 mg L⁻¹ PE).

97

98 *2.2 Cell growth*

99 At two-day intervals, air stones and stirrers were turned off, allowing PE particles to float to the
100 surface. A 2-mL aliquot from the central part of each cylinder was then pipetted into a cuvette and
101 absorption, as an optical density measure of cell growth (Choi and Lee, 2018), was determined at
102 565 nm with a UV-Vis spectrophotometer (Lambda 365, PerkinElmer). The dry mass of oven-dried

103 (60 °C for 24 h) pellet arising from the centrifugation (5 min at 5000 rpm) of a 4 mL aliquot sampled
104 concurrently was determined on a five-figure balance.

105 **2.3 End-points**

106 At the end of the four treatments, samples were vacuum-filtered through individual 25 µm nylon
107 filters (Hebei Reking) in order to capture remaining biomass and minimise contamination by PE
108 microplastics, and residues were oven-dried as above.

109 For the determination of chlorophyll a (Chl_a), chlorophyll b (Chl_b) and carotenoids, we used a
110 modified version of Arnon's method (Arnon, 1949). Briefly, 20 mg of dried microalgae were
111 extracted in 80% acetone and supernatants arising from subsequent centrifugation (as above) were
112 measured for absorption at 663 nm, 645 nm and 470 nm by UV-vis spectrophotometry.

113 Concentrations in µg mL⁻¹ were derived from the following formulae:

114 Chlorophyll a = $12.7(A_{663}) - 2.69(A_{645})$

115 Chlorophyll b = $22.9(A_{645}) - 4.68(A_{663})$

116 Total carotenoids = $[(1000(A_{470}) - 1.82 \text{ Chl}_a - 85.02 \text{ Chl}_b)] / 198$

117 The concentration of the proteinogenic imino acid, proline, that has a role in ameliorating
118 environmental stress in algae, was determined using acidic ninhydrin according to Bates et al.
119 (1973). The reagent was prepared by dissolving 0.625 g of ninhydrin in 25 mL of a solution consisting
120 of 15 mL glacial acetic acid and 10 mL of 6 M phosphoric acid and was stored at 4 °C. One hundred
121 mg of dried microalgae were ground with 10 mL of 3% sulphosalicylic acid in a porcelain pestle and
122 the resulting contents were centrifuged at 1500 rpm for 10 min. Two-mL aliquots of the extract,
123 ninhydrin reagent and glacial acetic acid were shaken in a foil-covered test tube and heated to 100
124 °C for 1 h in a water bath. The test tube was then cooled on ice before 4 mL of toluene were added
125 and the contents vortexed for 20 min before the absorbance of the upper solution was read at 520
126 nm by the UV-vis spectrophotometry after having been calibrated by proline standards in the range
127 of 0 to 40 mg mL⁻¹.

128 Malondialdehyde (MDA) was determined as an indicator of lipid peroxidation according to Haraguchi
129 et al. (1997). Thus, 50 mg of dried biomass were ground with 2 mL of 10% trichloroacetic acid in a
130 porcelain pestle and mortar and the resulting suspension was centrifuged at 10,000 rpm for 10 min.
131 One-mL of supernatant was added to 2 mL thiobarbituric acid and the contents heated in a bain-
132 marie for 45 min at 95 °C before the cooled suspension was centrifuged at 5000 rpm for 10 min. The

133 absorbance of the supernatant was read at 532 nm and corrected for nonspecific absorbance at 600
134 nm by UV-vis spectrophotometry.

135 **2.4. Microscopy and Raman spectroscopy**

136 Just before the termination of the experiment, 2-mL aliquots were pipetted from each container on
137 to individual microscope slides and the contents analysed under a Carl Zeiss binocular microscope at
138 up to 200 X magnification and an Olympus IX51 cell culture fluorescence microscope. Additional,
139 freeze-dried aliquots were analysed under a Tescan Vega 3 scanning electron microscope operated
140 at 20 kV and a viewing distance of between about 5 and 10 mm, and a μ -Raman spectrometer
141 (LabRAM HR, Horiba, Japan) employing a laser wavelength of 582 nm, a Raman shift of 400-1800 cm^{-1}
142 and acquisition times between 20 and 30 s.

143

144 **2.5. Statistics**

145 Mean values were compared by one-way ANOVA using SPSS. A Duncan's multiple range post-hoc
146 test used to identify statistical differences ($\alpha = 0.05$) between specific pairs of means.

147



148 **3. Results**

149 **3.1. Cell growth**

150 Cell growth of *A. platensis* is shown as a function of exposure time for the four different treatments
151 in Figure 1. Mean optical density (Figure 1a) is about 0.04 at the beginning of all treatment and, in
152 the control, increases to 1.0 at 20 h. After 12 h, mean optical densities in the presence of
153 microplastics PE are lower than the corresponding values for the control. However, the reduction is
154 only significant ($p < 0.05$), and in the range of about 10 to 15%, at the highest concentration of PE
155 employed.

156 Mean dry biomass (Figure 1b) is about 0.07 mg mL^{-1} at the beginning of all treatments and, in the
157 control, increases to about 1.2 mg mL^{-1} after 14 h. Increasing concentration of PE microplastics is
158 accompanied by a progressive reduction in mean dry weight after 6 h. Reduced dry weight is
159 significant relative to the control for microplastic concentrations of 10 mg L^{-1} and 100 mg L^{-1} after 8
160 h, and beyond 12 h dry weight is significantly lower at 100 mg L^{-1} than at 10 mg L^{-1} .

161




163

164 Figure 1: Cell growth of *A. platensis* as (a) optical density and (b) dry biomass as a function of
 165 exposure time for the four treatments. Errors are one standard deviation about the mean of four
 166 measurements.

167 Figure 2 shows the mean concentrations of chlorophyll a, chlorophyll b and total carotenoids in the
 168 biomass of *A. platensis* arising from the four exposures. Chlorophyll a exhibits a significant,
 169 progressive decrease with increasing concentration of PE microplastics, while chlorophyll b and
 170 carotenoids exhibit significant reductions relative to the corresponding controls are only observed at
 171 particle concentrations of 10 mg L⁻¹ and 100 mg L⁻¹. Overall, mean concentrations of chlorophyll a,
 172 chlorophyll b and total carotenoids decrease by about 70%, 50% and 40%, respectively, from the
 173 control to the highest concentration of microplastics added.

174




175

176 Figure 2: Concentrations of chlorophyll a, chlorophyll b and total carotenoids, normalised to dry
 177 weight of *A. platensis*, in the four treatments. Error bars represent one standard deviation about the
 178 mean ($n = 4$) and, for each component (and letter style), different letters indicate a significant
 179 differences ($p < 0.05$).

180

181 Figure 3 shows the mean proline concentration of *A. platensis* in the different treatments. Addition
 182 of 1 mg L⁻¹ of PE microplastics results in a proline concentration that is not significantly different to
 183 the control (about 7.5 $\mu\text{mol g}^{-1}$), but 10 mg PE L⁻¹ and 100 mg PE L⁻¹ result in progressive increases
 184 that are significant, with a mean proline concentration of about 32 $\mu\text{mol g}^{-1}$ in the highest exposure.

185



186

187 Figure 3: Concentrations of proline, normalised to dry weight of *A. platensis*, in the four treatments.

188 Error bars represent one standard deviation about the mean ($n = 4$) and different letters indicate

189 significant differences ($p < 0.05$).

190

191 Mean concentrations of MDA in *A. platensis* arising from the four treatments are shown in Figure 4.

192 Here, differences are smaller between treatments than for proline concentrations and significant

193 differences are only observed between the two highest exposures and the control and the lowest

194 exposure.


195

196

197

198

199



200

201 Figure 4: Concentrations of MDA, normalised to dry weight of *A. platensis*, in the four treatments.
 202 Error bars represent one standard deviation about the mean ($n = 4$) and different letters indicate
 203 significant differences ($p < 0.05$).


204

205

206 Figures 5 and 6 show binocular and fluorescent microscopic images, respectively, of *A. platensis*
 207 abstracted at the end of the experiment. Mature, helicoidal cells in the control reached lengths of
 208 **600 μm** and with diameters of about 5 μm , but increasing concentration of PE microplastics was
 209 accompanied by increasing fragmentation, with lengths often below 25 μm , swelling to about 8 μm ,
 210 and attachment of plastic. Also indicated in Figure 6 are microplastics attached or aggregated at
 211 regions where fragmentation has occurred or is about to take place. The attachment of microplastics
 212 to the algal surface more generally is evident when comparing the SEM images of the control and
 213 exposure to 100 mg PE L⁻¹ in Figure 7.

214


215



216


217 Figure 5: *A. platensis* under the binocular microscope in (a) the control, (b) exposed to 1 mg L⁻¹ PE,
218 (c) 10 mg L⁻¹ PE, (d) 100 mg L⁻¹ PE.

219



220

221 Figure 6: *A. platensis* under the fluorescent microscope in (a) the control, (b) exposed to 1 mg L⁻¹ PE,
 222 (c) 10 mg L⁻¹ PE, (d) 100 mg L⁻¹ PE. Arrows show fragmented or fragmenting regions where
 223 microplastics appear to be attached or aggregated.




224

225 Figure 7: SEM images of *A. platensis* in (a) the control and (b) exposed to 100 mg L⁻¹ PE.

226

227 Raman spectra between wave numbers of 900 and 1200 cm^{-1} are shown for *A. platensis* abstracted
228 at the end of the experiment and for the PE microplastics in Figure 8. The peaks centred at 1153 cm^{-1}
229 are related to carotenoid substances, and intensities are similar for the control and lowest
230 concentrations of microplastics but at the highest concentration of particles, and (at least
231 qualitatively) consistent with the observations in Figure 2, the peak is suppressed. The peak at about
232 at 1067 cm^{-1} is related to polyethylene (C-C stretching) and is clearest for the microplastics.
233 However, this peak is also evident in *A. platensis* exposed to microplastics and with an intensity that
234 decreases with microplastic concentration, confirming the algal-microplastic associations observed
235 microscopically above.

236



237

238 Figure 8: Raman spectra for *A. platensis* sampled at the end of the experiment and for the PE
239 microplastics.

240

241 4. Discussion

242 The results of this study show that PE microplastics do not have lethal effects on the cyanobacteria,
243 *A. platensis*, but are responsible for physiological and biochemical changes. Specifically, PE
244 microplastics inhibit the growth and reduce photosynthesis in *A. platensis* over the range of

245 concentrations studied (1 to 100 mg L⁻¹), with an EC-50 for growth on a dry mass basis of about 100
246 mg L⁻¹. Concentrations of 10 and 100 mg L⁻¹ also cause oxidative stress leading to lipid peroxidation,
247 and result in the intracellular production of proline.

248 In a recent review, Prato et al. (2019) highlighted the limited number of studies on the effects of
249 microplastics and nanoplastics on microalgae, and the lack of consensus among these studies. With
250 respect to growth, for example, the presence of microplastics have revealed inhibition (e.g. Besseling
251 et al., 2014; Mao et al., 2018), promotion (e.g. Lagarde et al., 2016; Canniff and Hoang, 2018) and no
252 effects (e.g. Davarpanah and Guilhermino, 2015; Yokota et al., 2017).

253 The growth inhibition ratio, IR, for *A. platensis* exposed to PE microplastics was calculated from
254 optical densities at 20 h in each exposure (E20) and control (C20) (Ansari et al., 2021):

$$255 \text{ IR (\%)} = (1 - E_{20}/C_{20}) \times 100\%$$

256 We note that ratios of 3.6%, 5.6% and 9.0% for PE concentrations of 1, 10 and 100 mg L⁻¹,
257 respectively, are lower than the typical range reported in the literature for different genera of
258 microalgae exposed to various polymers at concentrations between 50 and 250 mg L⁻¹ (about 20 to
259 50%, and where growth inhibition was observed; Besseling et al., 2014; Li et al., 2020; Ansari et al.,
260 2021). However, our results are, quantitatively, more consistent with growth rate inhibitions
261 reported for *Spirula* sp. exposed to microplastic concentrations above 25 mg L⁻¹ (Abed et al., 2021;
262 Hadiyanto et al., 2021; Hadiyanto et al., 2022). Presumably, discrepancies in the literature relate to
263 differences in species morphology and cell characteristics, as well as plastic particle size and surface
264 charge, the concentration and type of any additives present, and the exposure period employed
265 (Nava and Leoni, 2021).

266 Moderate growth rate suppression by microplastics in the present study was accompanied by
267 swelling of cells and fragmentation of filaments (Figures 5 and 6). The propagation of *A. platensis*
268 takes place by fragmentation of trichomes that are subsequently elongated by binary fission until
269 maturity (Jung et al., 2021). Increasing fragmentation in the presence of increasing quantities of MPs
270 may reflect a progressive delay of the propagation phase, or some response of mature microalga to
271 plastic. In a study of *A. platensis* exposed to cadmium, Rangsayatorn et al. (2001) attributed
272 fragmentation to a defence mechanism. Specifically, low concentrations (< 8 mg L⁻¹) of Cd resulted in
273 a small number of disordered cells, with higher concentrations resulting in severe injury.

274 Nevertheless, and despite cellular damage, growth and division continued. In the present study, the
275 attachment or aggregation of microplastics at the surface of the microalgae, evident in microscopic
276 imagery, may contribute to the weakening of filaments or trigger some defence. This may explain

277 why microplastic accumulation is concentrated at the cell terminal (Figure 6). Alternatively, it is
278 possible that MPs are attracted to the cytoplasm content that is released in this region during cell
279 fragmentation (Jung et al., 2021). The precise mechanisms behind this process and the potential for
280 small micro- or nanoplastics to be internalised during fragmentation are not known and warrant
281 further study.

282 As well as inhibiting the growth of *A. platensis*, exposure to PE microplastics affects the
283 photosynthetic activity of the microalga. Moreover, a reduction in the concentration of pigments
284 essential for photosynthesis was accompanied by increases in MDA content, as an indication of cell
285 membrane lipid peroxidation by reactive oxygen species (ROS), and proline, as an ROS scavenger and
286 antioxidant. Thus, ROS are formed when electrons are diverted to O₂ and not CO₂ when
287 photosynthetic activity decreases (Bhattacharya et al., 2010). Recent studies have also
288 demonstrated a reduction in pigment content of freshwater microalgae following exposure to
289 microplastics of different polymeric compositions at concentrations on the order of 100 mg L⁻¹ (Li et
290 al., 2020; Tunali et al., 2020). Potential causes of photosynthetic inhibition by microplastics include
291 shading from incident light (Zhang et al., 2017) and adherence to the algal surface via extracellular
292 polymeric substances, thereby reducing nutrient and gas exchange and trapping harmful
293 metabolites (Mao et al., 2018). It is not clear whether the reduction of photosynthetic activity is
294 related to or independent of growth inhibition and fragmentation in the presence of plastic but we
295 note that accumulation of ROS by *A. platensis* can result in spiral breakage by oxidizing lipids of the
296 sheath or cell membrane (Ma and Gao, 2010).

297 Although the effects observed here are broadly consistent with those reported elsewhere in the
298 literature, their ecological relevance is likely to be low because unrealistically high concentrations of
299 well-defined microplastics are typically considered in the presence of a single microalgal species.
300 Nevertheless, relatively small disruptions at the population level, including subtle changes to
301 morphology and palatability, could have more serious implications for ecosystem services and
302 functioning (Prata et al., 2019; Nava and Leoni, 2021). More specifically, and given its importance as
303 a human source of human protein (Habib et al., 2008), population-level alterations to *A. platensis*
304 could have significant impacts on food security. Conversely, it has been suggested that *A. platensis*
305 could be exploited to assist in the degradation of microplastics (Hadiyanto et al., 2021) or act as a
306 bioremediator through hetero-aggregation (Abed et al., 2021).

307

308 **5. Conclusions**

309 The present study has shown that exposure of the cyanobacterium, *A. platensis*, to different
310 concentrations of PE microplastics over a 20-d period results in various adverse effects, including a
311 reduction in growth rate and photosynthesis, oxidative stress, and the fragmentation and swelling of
312 trichomes. Although these impacts were observed for environmentally unrealistic concentrations of
313 well-defined, customised microplastics, small disruptions arising from lower concentrations could
314 have adverse effects at the population level. Because of the potential implications for ecosystem
315 services, further research into the impacts and interactions of lower concentrations of more
316 representative and heterogenous microplastics (in terms of size, polymer composition, surface charge
317 and degree of weathering, for example) is warranted.

318

319 **Acknowledgements**

320 We thank Shiraz University for financially supporting this study (grant number OGCB1M201526).

321

322 **References**

323 Abed, S.A., Kadhim, R., Alasady, A., Jazie, A.A. Materials Today@: Proceedings 42, 2251-2258. Ansari,
324 F., Ratha, S., Renuka, N., Ramanna, L., Gupta, S., Rawat, I., Bux, F., 2021. Effect of microplastics on
325 growth and biochemical composition of microalga *Acutodesmus obliquus*. Algal Research 56,
326 102296.

327 Arnon, D., 1949. Copper enzymes in isolated chloroplasts polyphenoloxidase in *Beta Vulgaris*. Plant
328 Physiology 24, pp. 1-15.

329 Bates, L.S., Waldren, R.P., Teare, I., 1973. Rapid determination of free proline for water-stress
330 studies. Plant and soil 39, 205-207.

331 Bhattacharya, P., Lin, S., Turner, J.P., Ke, P.C., 2010. Physical adsorption of charged microplastic
332 nanoparticles affects algal photosynthesis. Journal of Physical Chemistry 114, 16556-16561.

333 Canniff, P.M., Hoang, T.C., 2018. Microplastic ingestion by *Daphnia magna* and its enhancement on
334 algal growth. Science of the Total Environment 633, 500-507.

335 Choi, W.Y., Lee, H.Y., 2018. Enhancement of neuroprotective effects of *Spirulina platensis* extract
336 from a high-pressure homogenization process. Applied Sciences 8, 634.

337 Davarpanah, E., Guilhermino, L., 2015. Single and combined effects of microplastics and copper on
338 the population growth of the marine microalgae *Tetraselmis chuii*. *Estuarine and Coastal Shelf*
339 *Science* 167, 269–275.

340 Gigova, L., Marinova, G., 2016. Significance of microalgae – Grounds and areas. *Genetics and Plant*
341 *Physiology* 6, 85-100.

342 Habib, M.A.B., Parvin, M., Huntington, T.C., Hasan, M.R., 2008. A review on culture, production and
343 use of *Spirulina* as food for humans and feeds for domestic animals and fish. *FAO Fisheries and*
344 *Aquaculture Circular No. 1034*, Food and Agriculture Organization of the United Nations, Rome.

345 Hadiyanto, H., Khoironi, A., Dianratri, I., Suherman, S., Muhammad, F., Vaidyanathan, S., 2021.
346 Interactions between polyethylene and polypropylene microplastics and *Spirulina* sp. microalgae in
347 aquatic systems. *Heliyon* 7, e07676.

348 Hadiyanto, H., Muslihuddin, M., Khoironi, A., Pratiwi, W.Z., Fadilah, M.N., Muhammad, F., Afiati, N.,
349 Dianratri, I., 2022. The effect of salinity on the interaction between microplastic polyethylene
350 terephthalate (PET) and microalgae *Spirulina* sp. *Environmental Science and Pollution Research* 29,
351 7877-7887.

352 Haraguchi, H., Ishikawa, H., Kubo, I., 1997. Antioxidative action of diterpenoids from *Podocarpus*
353 *nagi*. *Planta Med.* 63, 213-215.

354 Jung, C.H.G., Braune, S., Waldeck, P., Küpper, J.H., Petrick, I., Jung, F., 2021. Morphology and growth
355 of *Arthrospira platensis* during cultivation in a flat-type bioreactor. *Life* 11, 536.

356 Lagarde, F., Olivier, O., Zanella, M., Daniel, P., Hiard, S., Caruso, A., 2016. Microplastic interactions
357 with freshwater microalgae: hetero-aggregation and changes in plastic density appear strongly
358 dependent on polymer type. *Environmental Pollution* 215, 331-339.

359 Li, S., Wang, P., Zhang, C., Zhou, X., Yin, Z., Hu, T., Hu, D., Liu, C., Zhu, L., 2020. Influence of
360 polystyrene microplastics on the growth, photosynthetic efficiency and aggregation of freshwater
361 microalgae *Chlamydomonas reinhardtii*. *Science of the Total Environment* 714, 136767.

362 Ma, Z., Gao, K., 2010. Spiral breakage and photoinhibition of *Arthrospira platensis* (Cyanophyta)
363 caused by accumulation of reactive oxygen species under solar radiation. *Environmental Experimentl*
364 *Botany* 68, 208–213.

365 Mao, Y., Ai, H., Chen, Y., Zhang, Z., Zeng, P., Kang, L., Li, W., Gu, W., He, Q., Li, H., 2018.
366 Phytoplankton response to polystyrene microplastics: Perspective from an entire growth period.
367 *Chemosphere* 208, 59-68.

368 Mobin, S., Alam, F., 2017. Some promising microalgal species for commercial applications: A review.
369 Energy Procedia 110, 510-517.

370 Nava, V., Leoni, B., 2021. A critical review of interactions between microplastics, microalgae and
371 aquatic ecosystem function. Water Research 188, 116476.

372 Prata, J.C., da Costa, J.P., Lopes, I., Duarte, A.C., Rocha-Santos, T., 2019. Effects of microplastics on
373 microalgae populations: A critical review. Science of the Total Environment 665, 400-405.

374 Priyadarshani, I., Rath, B., 2012. Commercial and industrial applications of micro algae – A review.
375 Journal of Algal Biomass Utilization 3, 89-100.

376 Rangsaytorn, N., Upatham, E.S., Kruatrachue, M., Pokethitiyook, P., Lanza, G.R., 2002.
377 Phytoremediation potential of *Spirulina (Arthrospira) platensis*: biosorption and toxicity studies of
378 cadmium. Environmental Pollution 119, 45-53.

379 Skjånes, K., Rebours, C., Lindblad, P., 2013. Potential for green microalgae to produce hydrogen,
380 pharmaceuticals and other high value products in a combined process. Critical Reviews in
381 Biotechnology 33, 172-215.

382 Tunali, M., Uzoefuna, E.N., Tunali, M.M., Yenigun, O., 2020. Effect of microplastics and microplastic-
383 metal combinations on growth and chlorophyll a concentration of *Chlorella vulgaris*. Science of the
384 Total Environment 743, 140479.

385 Yokota, K., Waterfield, H., Hastings, C., Davidson, E., Kwietniewski, E., Wells, B., 2017. Finding the
386 missing piece of the aquatic plastic pollution puzzle: interaction between primary producers and
387 microplastics. Limnology and Oceanography Letters 2, 91–104.

388 Zhang, C., Chen, X., Wang, J., Tan, L., 2016. Toxic effects of microplastic on marine microalgae
389 *Skeletonema costatum*: Interactions between microplastic and algae. Environmental Pollution 220,
390 1282-1288.

391

392