



PEARL

Degradation of Cornish calcified seaweed (maerl) in soil, and effects on soil fertility

Parkinson, R; Hall-Spencer, JM; Burchett, S; Young, H

Published in:

Proceedings of the Eurosoil Conference, Freiburg, Germany. 4-12 September 2004.

Publication date:

2004

Link:

[Link to publication in PEARL](#)

Citation for published version (APA):

Parkinson, R., Hall-Spencer, JM., Burchett, S., & Young, H. (2004). Degradation of Cornish calcified seaweed (maerl) in soil, and effects on soil fertility. *Proceedings of the Eurosoil Conference, Freiburg, Germany. 4-12 September 2004.*, 0(0).

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Degradation of Cornish calcified seaweed (maerl) in soil, and effects on soil fertility

Rob Parkinson¹, Jason Hall-Spencer², Sarah Burchett³, Helen Young⁴
School of Biological Sciences, University of Plymouth, Drake Circus, Plymouth, UK
1rparkinson@plymouth.ac.uk
2jhall-spencer@plymouth.ac.uk
3sburchett@plymouth.ac.uk
4hyoung@students.plymouth.ac.uk

Calcified seaweed (maerl) occurs extensively around the westernmost coasts of Europe, predominantly in the low intertidal zones offshore Spain, North West France, South West England and Western Ireland (Hall-Spencer et al., 2003). This material has been used extensively in agriculture, both as a general soil conditioner and also specifically as a liming agent and to supply trace elements (Tye et al., 2000)

The research reported here will discuss the degradation of calcified seaweed in soil, and consequent soil fertility benefits, notably soil respiration rates, associated with the use of Cornish Maerl (calcified seaweed) as an agricultural liming agent. Specifically, the following will be discussed:

1. Current extraction and utilisation of Cornish calcified seaweed
2. Scanning electron photomicrograph analysis of calcified seaweed microstructure following degradation in soil
3. The effects of maerl on soil aggregate stability, soil respiration rates and soil pH/extractable calcium concentrations.
4. Ryegrass response to calcified seaweed additions.

References

Hall-Spencer, J. M. Grall, J. Moore, P. G. & Atkinson, R. J. A. 2003. Bivalve fishing and maerl-bed conservation in France and the UK - retrospective and prospect. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13, S33-S41.

Tye, A., Fullen, M. & Hocking, T. 2000. Response of Grass Leys to Applications of Calcified Seaweed. *Communications in Soil Science and Plant Analysis* 31, (3 & 4), 529-542.