



PEARL

**A Pilot RCT Investigating the Effects of Targeted Compression on Athletes with Pelvic / Groin Pain**

Sawle, Leanne; Freeman, Jennifer; Marsden, Jonathan

**Published in:**

Journal of Sport Rehabilitation

**DOI:**

[10.1123/jsr.2017-0097](https://doi.org/10.1123/jsr.2017-0097)

**Publication date:**

2017

**Link:**

[Link to publication in PEARL](#)

**Citation for published version (APA):**

Sawle, L., Freeman, J., & Marsden, J. (2017). A Pilot RCT Investigating the Effects of Targeted Compression on Athletes with Pelvic / Groin Pain. *Journal of Sport Rehabilitation*, 0(0), 1-34. <https://doi.org/10.1123/jsr.2017-0097>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

1 **Accepted for publication in the Journal of Sports Rehabilitation. Acceptance date**  
2 **3/10/2017. <https://journals.humankinetics.com/doi/10.1123/jsr.2017-0097>**

3

4 **A Pilot RCT Investigating the Effects of Targeted Compression on Athletes with Pelvic /**  
5 **Groin Pain**

6 Leanne Sawle, Jennifer Freeman, Jonathan Marsden.

7 **AFFILIATIONS:** <sup>1</sup>Plymouth University, Faculty of Health and Human Sciences, Peninsula Allied  
8 Health Centre, Plymouth, Devon, United Kingdom. <sup>2</sup>Cardiff University, School of Engineering,  
9 The Parade, Cardiff, Wales.

10

11 **Context:** Athletic pelvic/groin pain is a common yet often challenging problem to both  
12 diagnose and manage. A new tool has been developed based on the clinical effects of applied  
13 force on the pelvis. Early findings indicate that this customised compression orthosis may  
14 have a positive effect upon pelvic/groin pain and performance measures.

15 **Objectives:** To:

16 Inform the design and test the practicality of procedures for a future definitively powered  
17 randomized controlled trial;

18 Provide an estimate of the effect size of this orthosis on selected clinical and performance  
19 measures.

20 **Design:** Pilot randomised controlled trial with participants randomly allocated to an  
21 intervention or waiting-list control group

22 **Setting:** The training location of each athlete

23 **Participants:** 24 athletes with sub-acute and chronic pelvic conditions were proposed to be  
24 recruited

25 **Intervention:** A customised compression orthosis, delivering targeted compression to the  
26 pelvic girdle.

27 **Outcome measures:** Measures were the active straight leg raise test, squeeze test, broad  
28 jump, and the multiple single-leg hop-stabilization test.

29 **Results:** 16 athletes completed the study. The invention group demonstrated moderate to  
30 large estimated effect sizes on the squeeze test and active straight leg raise tests ( $d = 0.6-1.1$ )  
31 whilst wearing the orthosis. Small effect sizes ( $d = 0.2$ ) were seen on jump distance and the  
32 dominant leg balance score. Compared to the control group the intervention group also  
33 showed moderate to large estimated effect sizes on the active straight leg raise measures ( $d =$   
34  $0.5-0.9$ ) when wearing sports shorts.

35 **Conclusions:** The protocol was feasible. Effect sizes and recruitment/attrition rates suggest  
36 that the intervention holds promise and that a future definitive powered RCT appears feasible  
37 and is indicated.

38

39

40 **INTRODUCTION**

41

42 The incidence of pelvic/ groin injury is particularly high in sports such as Gaelic (24%)<sup>1</sup>, ice  
43 hockey (10-11%)<sup>2</sup> and Association Football (49%)<sup>3</sup>, and research has highlighted the  
44 challenges affecting the diagnosis and management of these injuries <sup>4-6</sup>.

45 Pelvic belts, a form of external pelvic compression <sup>7</sup>, are a tool that have demonstrated some  
46 success in reducing pain and improving function, on clinical tests such as the squeeze test and  
47 active straight leg raise (ASLR) <sup>8,9</sup>. However, the practicality of using belts during  
48 performance is limited, and research has begun to consider alternative forms of external  
49 pelvic compression. Preliminary research has suggested that pain and/ or function on clinical  
50 tests (ASLR and squeeze test force) may be improved by introducing targeted compression in  
51 the form of a customised compression orthosis. Subjective data from this study further  
52 proposed positive effects upon attributes including power and balance<sup>10</sup>. It has been  
53 hypothesised that these effects may be explained by targeted compression influencing the  
54 force or form closure deficit associated with this type of injury (providing stability), and/or  
55 improving proprioception and muscle function<sup>10</sup>.

56 The use of compression garments as a post exercise adjunct to recovery, have been reported  
57 as beneficial for performance recovery and delayed-onset muscle soreness<sup>11,12</sup>. However,  
58 there is a paucity of research in the field of compression and injury management. Of the work  
59 undertaken in this domain, one study reported that standard compression shorts have been  
60 found to significantly reduce pain in athletes with osteitis pubis<sup>13</sup>. Other work found that  
61 targeted compression reduced adductor activity in healthy participants, and theorised as  
62 reducing the risk of adductor related injury<sup>14</sup>.

63 Research into compression style orthoses has reported mixed findings in terms of enhancing  
64 performance attributes; some studies demonstrating improvements in measures such as  
65 balance and power, whilst others showing no effect<sup>13,15,16</sup>. Some findings have suggested that  
66 compression shorts may influence repetitive performance by reducing muscle oscillations<sup>15</sup>,  
67 influencing proprioception and delivering athlete perceived improvement effects<sup>16</sup>. Well-  
68 fitting compression shorts have demonstrated improvements in the static balance of female  
69 athletes, compared to wearing standard shorts<sup>17</sup>

70 However, little is known about the application of targeted compression, and this warrants  
71 further investigation. Whilst acknowledging the limited literature in this domain, there is  
72 some evidence that targeted compression may have a role in athletic groin and pelvic injury  
73 management<sup>13,14</sup>. It is also possible that a customised targeted compression orthosis, may  
74 offer further benefits.

75 Therefore, to explore the role of compression in injury management, and specifically the use  
76 of external pelvic compression in the form of a customised compression orthosis, a  
77 randomized controlled trial (RCT) was indicated. However several factors must be  
78 determined prior to designing and implementing a full trial, therefore a pilot RCT<sup>18</sup> was  
79 undertaken in order to:

80 Inform the design and test the practicality of procedures for a future definitively powered  
81 RCT study<sup>19</sup>, by determining:

- 82 a. recruitment rate
- 83 b. attrition rate
- 84 c. presence of adverse events
- 85 d. effect size estimate
- 86 e. feasibility of using the outcome measures

87 f. effectiveness of the blinding strategy

88 g. practicality of the protocol

89

90

## 91 **METHOD**

92

### 93 **Sampling and Recruitment Strategy**

94 A convenience sample of volunteers was recruited from UK-based sports clubs over one year.

95 The number of males recruited may reflect the fact that that moderate evidence exists

96 supporting a higher risk of this type of injury in male athletes <sup>20</sup>.

97 Table 1 presents the demographical data.

98 Of the nine athletes allocated to the waiting-list control group, eight had chronic pain; one

99 athlete was identified as having sub-acute pain during screening, but this became chronic pain

100 (> three months) by the time the baseline measures were taken. In the intervention group, all

101 seven athletes had chronic pain. Pain severity ranged from low to moderate across both

102 groups and was influenced by activity; as per the exclusion criteria those exhibiting severe

103 pain (>8/10 on a numerical rating scale [NRS]) did not take part. This was for ethical reasons

104 due to repeated testing.

105 The uneven numbers in the two groups were due to the minimisation program which was

106 setup for 12 athletes in each group; split evenly across chronic and sub-acute pain.

107

108

109 **Table 1 Athlete demographics**

110 All athletes were training three or more times per week, and were undertaking both aerobic  
111 and anaerobic training. Competition levels ranged from recreational (n = 8) to professional (n  
112 = 8).

113

## 114 **Eligibility Criteria**

### 115 *Inclusion Criteria:*

- 116 i. Athletes aged 18 years or above (recreational or professional).
- 117 ii. Sub-acute (1-3 months duration) and chronic (>3 months) self-reported pelvic / groin  
118 pain presenting during sport or at rest (unilateral or bilateral)
- 119 iii. Pelvic / groin pain as confirmed via a screening procedure. For inclusion, positive pain  
120 scores had to be determined on at least two of these five tests, as when used in isolation these  
121 tests are limited in terms of reliability, but when used together they provide a more reliable  
122 approach <sup>21</sup>.

123

## 124 **Screening Procedure**

125

126 The following battery of tests were performed; details can be found in previous literature.

127 These tests are appropriate for both unilateral and bilateral pain presentations<sup>8,21,22</sup>:

- 128 • Active Straight Leg Raise (ASLR)
- 129 • Fabers
- 130 • Thigh thrust
- 131 • Gaenslens
- 132 • Squeeze test

133

134 *Exclusion Criteria:*

135 Self-reported acute pelvic / groin pain; defined as zero to four-weeks duration, which may be  
136 expected have a short resolution period <sup>23</sup>.

137 Neurological, or systemic disease

138 Pregnancy

139 Radicular pain

140 History of pelvic fracture

141 Inguinal hernia

142 Severe pain (>8/10 on a NRS)

143 Trochanteric bursitis

144 Ruptured muscle

145

146 **Study Design**

147 A waiting-list control <sup>24</sup>, researcher blinded <sup>25</sup>, pilot RCT was undertaken after approval from  
148 the local (UK-based University) ethics committee. A waiting-list control design was  
149 employed for ethical reasons, as all athletes were selected on the premise that they were  
150 suffering from ongoing pain. This is considered a useful method for keeping the control  
151 athletes engaged with the study<sup>26</sup>.

152 Random allocation with a minimisation procedure was employed to ensure equal distribution  
153 of sub-acute and chronic conditions between groups. Athletes in the intervention group used



154 the compression orthosis for a four-week period. Athletes in the control group served as a  
155 waiting-list control for a four-week period, before receiving their orthosis.

### 156 **Recruitment Rate**

157 The recruitment and attrition rates were reported according to CONSORT Guidelines <sup>27</sup>.

### 158 **Sample Size**

159 Twenty-four athletes were proposed to be randomly assigned to the intervention (n=12) or  
160 waiting-list control group (n=12), based on the recommendation of 12 in each athlete group  
161 for feasibility/pilot work <sup>28</sup>.

162

163 Figure 1 summarises the athletes' route through the study.

164

### 165 **Figure 1 Athlete pathway through the study**

166

167 After obtaining written informed consent, potential athlete participants were screened, and  
168 demographic, pelvic /groin pain history and training data (frequency, duration and type) were  
169 recorded. Athletes who met the eligibility criteria were measured for a compression orthosis  
170 by the investigator.

171 One week later (+/- 3 days for flexibility) athletes completed two sets of baseline measures  
172 wearing sports shorts and loose-fitting track pants over the top (provided). The use of track  
173 pants was to standardise dress, and to ensure blinding of the investigator at later dates.

174 Athletes were also fitted for their customised orthosis, and given usage and care instructions.  
175 The study administrator's details were provided for any future compression orthosis queries,  
176 and the orthosis held by the administrator until after the randomisation process.

### 177 **Randomisation Procedure**

178  
179 The administrator randomly allocated the athlete to the groups using a web-based system  
180 (minim <http://www-users.york.ac.uk/~mb55/guide/minim.htm>). A minimisation algorithm  
181 was used to ensure balance between the groups on the basis of injury chronicity (1-3 months  
182 versus > 3 months). Allocation concealment was employed to blind the investigator to the  
183 randomisation process <sup>29</sup>. The administrator informed athletes of their group allocation,  
184 posted the diaries to record training frequency, duration and type, treatment and compression  
185 orthosis usage, and sent the compression orthosis to the intervention group.

### 186 **Allocation Concealment during Outcome Measurement**

187  
188 A compression orthosis may have an orthotic effect<sup>30</sup>, only seen when the orthosis is in situ.  
189 Long term use of the compression orthosis may also result in improvements in the outcome  
190 measures even when it is not worn; a "carryover effect". To measure these potential effects  
191 the intervention group were assessed with and without the compression orthosis. For the  
192 intervention group, one assessment was completed when wearing the compression orthosis  
193 and another with shorts. Athletes in the control group were assessed twice with shorts. As  
194 there is a potential order effect the order of the testing (orthosis versus shorts) was  
195 randomised to account for effects such as fatigue or exacerbation of symptoms with testing.  
196 The administrator randomised the wearing of the orthosis, completed paper slips recording  
197 this information, and sealed them in opaque envelopes labelled with the athlete's name, study  
198 number and the measurement session number. These were sent to the investigator prior to

199 each test date so that they could hand the sealed envelope to the participant at the start of each  
200 session. Envelopes were also prepared for those in the waiting-list control group; the contents  
201 asked these athletes to wear shorts for both assessments. Athletes were asked to verbally  
202 confirm that the envelope that they had been given was sealed and had their name on it and  
203 the measurement session (week two, four or six) via a digital recorder.

## 204 **Blinding**

205

206 A criticism regarding the reporting of blinding in studies, is that many studies do not test the  
207 effectiveness of their blinding strategy<sup>25</sup>. Therefore, athletes wore track pants to conceal  
208 what they were wearing, and at week two, photographs were taken of athletes from the torso  
209 down at the start of assessment one and assessment two. To determine the effectiveness of the  
210 investigator blinding procedure, at the end of the study eight individuals were independently  
211 asked to identify whether a participant was wearing a compression orthosis or not from  
212 looking at the photographs. Further, at the end of the measurement sessions at week two, four  
213 and six the investigator completed a form indicating what they felt the athlete was wearing.

## 214 **Groups**

215

### 216 *Intervention Group*

217 Athletes were asked to wear their orthosis for normal training/ sport / physiotherapy input for  
218 a four-week “intervention” period and complete daily diaries to record usage, training, sport  
219 and physiotherapy input throughout this period.

### 220 *Waiting-List Control Group*

221 Athletes were asked to continue normal training/ performance/ physiotherapy input and  
222 record this in their daily diaries for a four-week period. After this time, the control group  
223 received the orthosis by post from the administrator.

## 224 **Timing and Purpose of Assessments**

225

226 Outcome measures were recorded at week one (baseline), week two, week four and week six,  
227 and athletes were assessed twice (assessment one and assessment two), separated by 10  
228 minutes of rest. The outcome measures were undertaken in a standardised order, and  
229 performed as described below.

230 The measures taken at baseline, when all athletes wore sports shorts for both assessments  
231 give an indication of the stability of the outcome measures over time. This was checked using  
232 intraclass correlation coefficients (ICCs) and Bland Altman plots.

233 To maximise recruitment testing was conducted in the athletes' clubs/sports centres using  
234 portable equipment to fit in around the athlete's schedule. To minimise the effects of external  
235 cues such as audience and environmental effects, athletes were tested in the same  
236 environment with only the investigator present.

## 237 **Outcome Measures**

### 238 *Primary Outcome Measure*

239 **Squeeze test** – Athletes with longstanding groin pain have shown significantly ( $p = <0.01$ )  
240 lower squeeze test force values than healthy controls<sup>31</sup>. This suggests that this test is  
241 appropriate for measuring the deficits associated with this type of pain. It has also shown  
242 excellent inter and intra tester reliability in athletes with and without groin pain (ICC  
243  $\geq 0.90$ )<sup>32</sup>.

244 From a supine position (hips and knees at 0°) athletes were asked to squeeze their legs  
245 together as hard as possible. This position has shown higher force output <sup>33</sup>, and minimal  
246 variability <sup>34</sup>. Maximal force output was measured using a padded load cell (SGA Applied  
247 Weighing, Reading, UK) placed between the medial femoral condyles, an oscilloscope (HPSI  
248 40i handheld pocket scope, Velleman Instruments, Taiwan) and an amplifier (Applied  
249 Weighing, Reading, UK). The voltage recorded was converted into Newtons.

#### 250 *Secondary Outcome Measures*

251 A familiarisation session was built into the start of the baseline testing session, so that  
252 athletes became aware of how to complete each test. Athletes had the tests verbally explained  
253 to them, could view the tests, look at photographs of the tests being performed, read simple  
254 instructions, and practice once on each leg.

#### 255 *The Active Straight Leg Raise (ASLR)*

256 Previous findings showed that the ASLR test produced low pain scores in a similar sample of  
257 athletes <sup>9</sup>, therefore the original ASLR protocol which records difficulty in completing the  
258 test <sup>35</sup> was also used. Research has also indicated that increased difficulty with the ASLR is  
259 reflected in higher pain scores on the test <sup>36</sup>.

260 In terms of reliability, its test retest reliability in post-partum posterior pelvic pain patients is  
261 excellent (ICC 0.87). Although reliability values are not available for athletes, the test has  
262 been used with athletes (from a variety of team and individual based sports) with groin pain  
263 <sup>8,37</sup>.

264 From a supine position on a plinth, athletes were asked to raise their right leg (knee in  
265 extension) to a bar positioned 20cm above the plinth. Athletes were asked to rate their pain at  
266 completion of the test using a numerical rating scale (NRS) of zero to ten (zero = no pain, ten  
267 = worst pain imaginable). Athletes were also asked to self-score the difficulty of this task

268 using a rating of zero to five (zero = no difficulty; five = extremely difficult). This was  
269 repeated with the left leg.

### 270 ***The Broad Jump***

271 The broad jump test of power<sup>38</sup> has been reported as demonstrating excellent test re-test  
272 reliability (ICC = 0.97)<sup>39</sup>. Athletes were asked to jump forwards over a mat, taking off from  
273 a two-footed stance and using their arms to propel themselves forward, landing with their feet  
274 close together. The landing spot was recorded using a chalked mark, and a right-angled tool  
275 (a hinged wooden bar) was used to measure from the landing mark, to the measuring tape  
276 fixed to the length of the mat. The protocol described by Almuzaini and Fleck<sup>39</sup> was  
277 followed, and the jump was repeated three times with the furthest distance recorded as their  
278 score.

### 279 ***Functional Balance***

280 The Multiple Single-Leg Hop-Stabilisation test (MSLHST) has been used as a functional,  
281 dynamic measure of athletic balance<sup>40</sup>, as due to its forward, transverse and diagonal  
282 movements, it tests balance across multi-movement planes. It has demonstrated good to  
283 excellent test re-test reliability in an active population (ICC = 0.85; CI 0.61-0.95)<sup>41</sup>.

284 Athletes were asked to jump from a standardised unipedal stance to and from 10 squares  
285 placed at distances determined by their height. The test incorporated periods of landing and  
286 statically maintaining a unipedal stance, giving athletes a balance and landing score for each  
287 of the 10 squares. The protocol reported by Riemann et al.<sup>40</sup> was used, and the test was  
288 undertaken on both the dominant and non-dominant leg. Leg dominance was defined as the  
289 leg with which the athlete prefers to kick with<sup>40</sup>, usually the right leg, therefore the left leg  
290 takes a pivotal role in providing stability<sup>42</sup>.

291 **Statistical Analysis**

292 Results were reported according to CONSORT Guidelines<sup>27</sup>.

293 To establish whether outcome measure scores could be averaged at baseline and for the two  
294 assessments per measurement session taken by the waiting-list control group, test retest  
295 reliability was examined where it was not already known in this patient group. ICCs (2,1)  
296 and Bland-Altman plots were used.

297 Fisher's Exact test was used to assess the effectiveness of the blinding procedures. It is a test  
298 used to analyse 2 x 2 contingency tables, and is advised for use with small sample sizes<sup>43</sup>.

299 Blinding is considered effective if no significant difference is seen between the responses  
300 given (incorrect and correct; p= 0.05).

301 Descriptive statistics were used, as recommended for pilot studies where a powered sample  
302 has not been employed<sup>44</sup>. Effect sizes were calculated (Cohen's d) and interpreted as being  
303 small =  $\geq 0.2$ ,  $<0.5$ , medium =  $\geq 0.5$  or large =  $\geq 0.8$ <sup>45</sup>. The formula for calculating effect  
304 sizes using Cohen's d is shown below (M = mean; SD = standard deviation).

$$d = \frac{M_1 - M_2}{SD_{\text{pooled}}}$$

305

306 An intention-to-treat approach to the descriptive analysis was employed in order to include  
307 data from all athletes randomized to a group, ignoring anything that occurs post  
308 randomisation<sup>46</sup>. The last measure carried forward technique was used in order to deal with  
309 any missing data from athletes dropping out during the study, and provided a conservative  
310 estimate of their performance had they continued<sup>47</sup>.

311 **Criteria to proceed to full RCT**

312 In order to determine feasibility<sup>19</sup>, the following criteria was required:

- 313 1. The attrition rate is <20% across the length of the study<sup>48</sup>.
- 314 2. The proposed number of athletes (n=24) could be recruited over a 12-month period.

315

316 **RESULTS**

317

318 The CONSORT diagram (figure 2) shows the numbers of athletes recruited, allocated to each  
319 group, and completing the study. **T tests showed no significant difference between the groups**  
320 **in terms of age, training, height or weight (P = >0.05).**

321

322

323 **Figure 2 The CONSORT diagram showing the flow of athletes through the study**

324

325 *Reliability*

326 ICCs and Bland Altman plots indicated that it was appropriate to average the waiting-list  
327 control group measures, and the baseline measures for both groups across assessment 1 and 2.  
328 This decision was justified by the ASLR ICC values indicating good to excellent reliability  
329 and precision (0.90-0.96; CI = 0.73-0.98). Bland Altman plots showed that the majority of the  
330 difference in test retest values stayed within 2SD. The decision to average the other outcome  
331 measures was based upon their historical intra-rater reliability.

332 *Effect Sizes*

333 Table 2 presents Cohen's d effect sizes representing the standardised mean difference in the  
334 scores of the intervention group compared to the waiting-list control group, at each stage of



335 the study. Table 3 shows descriptive statistics i.e. standardised mean differences and 95%  
336 confidence intervals.

337

338 **Table 2 The effect sizes (d) for each outcome measure at each stage of the study**

339

340 **Table 3 The mean difference (in bold) and 95% confidence intervals for the mean**  
341 **difference, from baseline to assessment week two, four and six for each outcome**  
342 **measure and for each condition**

343

344

345 *Blinding*

346 Eight individuals were asked to decide whether participants were wearing a compression  
347 orthosis or not by looking at photographs taken during the week two measurements. The  
348 responses were grouped as being either correct or incorrect. Fisher's Exact test indicated that  
349 there was no significant difference between the groups ( $p = 0.4$ ).

350 For the investigator's blinding check of effectiveness Fisher's Exact test showed that this  
351 result was not significant ( $p = 1$ ); blinding was effective.

352

## 353 **DISCUSSION**

354 Athletic pelvic/ groin pain is often a challenge both diagnostically, and from a management  
355 perspective<sup>5</sup>. Findings have suggested that the use of external pelvic compression in the form  
356 of a customised compression orthosis, may offer a tool for supporting the multi-modal  
357 management of these injuries<sup>10</sup>. However before implementing a full trial, a pilot RCT<sup>18</sup> was  
358 needed to inform the design and test the practicality of procedures for a future definitively  
359 powered RCT study<sup>19</sup>. These findings have been reviewed.

360 *Recruitment and Attrition Rates*

361 Of the intended 24 athletes, only 16 athletes (males = 13) were randomly assigned to groups  
362 and tested. Although the CONSORT diagram highlights the problem of ineligibility, it does  
363 not show that another 11 information packs were requested and received by interested athlete  
364 participants. This suggested that sufficient numbers were available, but that the 12-month  
365 study duration may have been an issue. Future work must consider time constraints, and how  
366 to improve the recruitment rate. Once recruited the attrition rate was zero demonstrating that  
367 once enrolled in the study, athletes were engaged enough to continue. It may also reflect that  
368 the attrition rate was not influenced by factors such as illness and other injuries.

369 *Adverse Effects*

370 No adverse effects were reported.

371 *Feasibility of Procedures and Outcome Measures*

372 Testing procedures proved to be successful in terms of logistics, practicality of outcome  
373 measures and data collection. The measures were straightforward to administer and athletes  
374 reported no difficulties in completing them. There was no missing data.

375 *Blinding Effectiveness*

376 The blinding procedures proved effective, and suggested that this method of blinding would  
377 be appropriate for future work.

378 *Summary of Outcome Measure Effect Sizes*

379 The results show that the compression orthosis had varying effects on a range of outcomes in  
380 athletes with chronic pelvic / groin pain. In general, wearing the compression orthosis  
381 demonstrated moderate to large effects on clinical measures, and negligible to small effects  
382 upon performance measures. These findings were considered and compared to the results of

383 compression studies, however the use of customised, targeted compression differs, and thus  
384 stands as a unique concept.

### 385 *Clinical Measures*

386 At week six, those allocated to the intervention demonstrated reduced pain and less difficulty  
387 in undertaking the ASLR, and, increased squeeze test force ( $d = 0.6$  to  $1.1$ ) compared to those  
388 in the waiting-list control group.

389 Moderate to large effect sizes ( $d = 0.5$  to  $0.8$ ) were seen on the ASLR measures when the  
390 intervention group were tested wearing sports shorts, indicating a possible carryover effect  
391 from orthosis use. The ASLR difficulty scores showed larger effect sizes ( $d = 1.1$ ) than pain  
392 on ASLR ( $d = 0.6$  to  $0.9$ ), supporting its appropriateness in this patient group<sup>8</sup>, and  
393 suggesting that other factors can influence performance difficulty. For example, increased  
394 pelvic mobility has been identified as a factor associated with higher ASLR scores<sup>35</sup>. This  
395 suggests those with more pelvic joint mobility find the ASLR test more difficult. In  
396 consensus with other research<sup>36</sup>, increased difficulty with the ASLR corresponded to higher  
397 pain scores on the test.

398 The large effect on squeeze test force ( $d = 0.8$ ) present at week four and week six, concurs  
399 with the effects of external pelvic compression on athletes with adduction-related groin pain  
400<sup>8</sup>. The findings from the intervention group wearing sports shorts indicates that this effect was  
401 associated only with wearing the orthosis. This may suggest a splinting or orthotic effect  
402 based on the use of an aid which demonstrates an effect only whilst it is in use. This could be  
403 explored with a longer intervention period to establish if a carryover effect becomes evident.  
404 Effects upon the ASLR support previous work in patients with chronic pelvic pain finding  
405 less ASLR difficulty with compression<sup>49</sup>. There is also support for the findings of  
406 compression orthoses reducing pain in athletes with osteitis pubis<sup>13</sup>. However, the

407 practicality of this compression orthosis, and its customised fit, may offer an improved  
408 method of applying targeted compression.

### 409 *Performance Measures*

410 Small effect sizes were seen on the broad jump and non-dominant leg MSLHST (d = 0.2 to  
411 0.3 respectively) when wearing the orthosis. A negligible effect was seen on the dominant leg  
412 (d=0.1).

413 Studies into compression shorts have shown contradictory findings on balance and power  
414 tests in healthy and patient populations. Whereas static unipedal balance has been seen to  
415 significantly improve (p = <0.05) when well-fitting compression shorts have been worn <sup>17</sup>,  
416 other findings demonstrated that compression shorts worn by healthy participants showed no  
417 significant effect (p = 0.9) upon static balance <sup>50</sup>. However, the static nature of this test may  
418 not have been athletically challenging or adequately responsive for a patient population.

419 In an athletic population with pelvic and groin pain, athletes with osteitis pubis showed a  
420 trend towards improved functional stability <sup>13</sup>; single leg squat (p = 0.08); effect size of d =  
421 0.2. This finding was for the left leg and may have indicated improved performance on the  
422 leg commonly required to provide stability for the dominant leg to perform. This finding  
423 might have partly explained the pilot RCT finding of a small effect seen in the non-dominant  
424 leg MSLHST score (d = 0.3), but minimal improvement seen on the dominant leg (d = 0.1).  
425 Due to bilateral pain reported in all athletes, and the ASLR mean differences and SD showing  
426 no difference between right and left leg pain scores, the effect of site of pain is unknown. It  
427 has been suggested that leg dominance should be considered in terms of the nature of the  
428 task, with the right leg dominating in activities requiring manipulation, for example kicking,  
429 whereas the left leg dominates in postural control activities <sup>42</sup>. The small improvement in the  
430 non-dominant or postural control leg, may have indicated that the targeted pelvic

431 compression led to small but identifiable improvements in the dynamic balance of athletes  
432 with pelvic / groin pain.

433 Field tests of power have also produced mixed findings in healthy athletes, from no  
434 significant effect of compression upon vertical jump height <sup>50</sup>, to customised compression  
435 shorts demonstrating significant improvements in countermovement vertical jump height (p =  
436 0.015)<sup>15</sup>. Whilst compression shorts did not improve maximal vertical jump power, a  
437 significant effect upon repeated jump performance was reported <sup>16</sup>. Mean power output on  
438 repeated jumps (n = 10) significantly improved when compression shorts were worn <sup>16</sup>.  
439 Compression leggings have also been shown to improve repeated sprint performance in  
440 healthy female athletes. Although there was no effect seen on haemodynamic or  
441 physiological measures, an influence upon proprioception was suggested <sup>51</sup>. This might have  
442 been due to the stimulation of the neuromuscular system. Gluteal muscle kinesiotaping has  
443 been found to increase explosive power as measured by a field test <sup>52</sup>.

444 This pilot RCT concurs with previous findings that wearing targeted compression shorts  
445 shows some improvement in power, but contributes new preliminary knowledge that this  
446 finding has been observed in athletes with pelvic / groin pain, and by using a customised  
447 approach.

#### 448 ***Intervention Assessment Points***

449 The effect sizes at different stages of the intervention period showed variable results. Week  
450 two improvements in the intervention group whilst wearing control shorts, may have  
451 indicated an immediate carryover. It is also possible that this was the influence of being  
452 allocated to the intervention group, and behaving accordingly.

453 However, the data varies over time; performance in the intervention group appears to have  
454 been detrimentally affected in the earlier assessment sessions before showing improvement at  
455 the latter assessments. One possible explanation may be that athletes underwent a period of  
456 adjustment to wearing the orthosis, and that there was variability in how they responded;  
457 possibly influenced by their expectations. It may also have been the result of increased  
458 discomfort caused by factors including the compression orthosis, increased training loads and  
459 changes in their condition. Early outcome measures may have been influenced by the level of  
460 pain at the start of the study, particularly in a population with varying pain mechanisms, and  
461 sites of pain <sup>4</sup>. Attempts were made to balance injury chronicity in both groups by way of a  
462 minimisation procedure. However, in view of the chronic nature of all participants, future  
463 work might wish to use a minimisation procedure to allocate athletes according to pain levels.  
464 Apprehension when undertaking measures for the first time may also have led to a tentative  
465 technique.

466 Although the performance measures showed small effect sizes, there may have been a  
467 learning effect, indicated by the control group also showing some improvements. Whilst  
468 effort was made to limit this by having a familiarisation session at baseline, balance studies  
469 have reported learning effects <sup>40,53</sup>. This may also have indicated an improvement in their  
470 condition.

#### 471 *Pain Provocation Tests*

472 Athlete responses to the five pain provocation tests ranged from two, to five positive  
473 outcomes. This figure was higher when bilateral pain responses are observed, and concurred  
474 with other studies finding bilateral and multiple sites of pain <sup>8</sup>. As none of the athletes  
475 presented with truly unilateral pain, the presence of bilateral pain might be indicative of those  
476 presenting with chronic pain. It is therefore not known if unilateral pain would have  
477 influenced the results.

478 As is expected with an inclusion criteria designed to identify athletes with any pelvic /groin  
479 pain; pain presentations varied in terms of the site(s) of pain <sup>4</sup>. It is also possible that this  
480 might have influenced the results, especially if positive pain responses were more evident in  
481 one group. However, there was an evenly matched spread in the number of positive pain  
482 responses across both groups. Therefore, it is suggested that this reflected the very nature of  
483 this injured population, and that pain presentations had a limited effect on the results. Despite  
484 this, future work should consider the number of positive pain responses as a minimisation  
485 factor.

#### 486 *Recruitment*

487 Future work requires an essential change to recruitment strategies. Sources of recruitment  
488 proved to be effective in generating interest from prospective athletes, but not in recruiting  
489 them into the study. This could have been due to the time commitment involved. Once the  
490 participant information pack was received 11 potential athletes were lost for reasons  
491 including work commitments. Of those recruited, ineligibility and the time/ resources  
492 available reduced the number of athletes completing the study. Having co-investigators may  
493 have helped, and been more efficient timewise when multiple athletes were being tested.

#### 494 *Limitations*

495 Although the intention was to recruit athletes with sub-acute and chronic pain, only the latter  
496 were recruited. Therefore, the results should be only considered in the context of a future  
497 study into chronic athletic pelvic / groin pain, and suggests it is more appropriate to focus  
498 upon recruiting athletes with chronic pain. **It is also noted that using a mixed sample of  
499 professional and recreational athletes is a confounding factor. The training / competition  
500 demands on the professional athletes may have influenced pain.**

501 This was also a partially blinded study which may have been influenced by demand  
502 characteristics <sup>54</sup>. Athletes knew which group they have been assigned to, and may have

503 adopted behaviour which they consider the investigator is demanding from them. This may  
504 have led to those wearing the compression orthosis trying hard to improve their performance  
505 to “please” the investigator. This may explain some of the positive effects seen at week two,  
506 when the compression orthosis was initially provided. At week two, even wearing shorts led  
507 to improvements in the intervention group (ASLR pain and difficulty scores). As the order of  
508 testing was randomised this cannot be explained fully by an instantaneous “carry-over  
509 effect,” as not all participants would have worn the orthosis first. Despite this possible bias,  
510 double-blinding was rejected because the effects of other compression shorts<sup>13,16</sup> would not  
511 allow for a true control. Therefore the reporting of blinding procedures was made transparent,  
512 and its effectiveness tested<sup>25</sup>.

### 513 *Contribution to Knowledge*

514 This pilot study has provided preliminary evidence to demonstrate the potential for  
515 employing a novel method of applying targeted compression to the pelvic girdle. Based on  
516 moderate to large improvement effects on clinical tests (ASLR and the squeeze test), and  
517 small improvement effects upon performance measures (balance and power), it is suggested  
518 that this unique compression orthosis may offer a practical tool to support the difficult  
519 management of chronic athletic pelvic / groin pain.

### 520 *Clinical Implications*

521 There may also be scope to explore the use of this compression orthosis in the prevention of  
522 pelvic / groin injury. Based on the findings of decreased adductor and biceps femoris activity  
523 with compression in both healthy and pelvic pain groups<sup>14,55</sup>, and the risks associated with  
524 increased and asymmetric activation, there may be a preventive role for this orthosis. As  
525 previous pelvic / groin injury is a risk factor for further injury<sup>56</sup>, this group of athletes would  
526 be appropriate to consider for orthotic use.



527 The possibility of a compression orthosis associated thermal effect upon performance should  
528 also be considered. Studies have reported that compared to control shorts compression shorts  
529 can significantly increase skin temperature during exercise (~1 degree centigrade)<sup>57</sup>, and that  
530 there is a relationship between increased skin temperature and increased muscle temperature  
531<sup>15</sup>. It has also been shown that during short duration exercise neuromuscular function can be  
532 affected by muscle temperature; functions such as nerve conduction velocity improving with  
533 higher temperatures. Improved performance has been also observed on vertical jump tests of  
534 power after the lower limbs have been heated<sup>58</sup>. This suggests that it may be appropriate to  
535 explore the use of this compression orthosis after warm up exercise, as this may show  
536 different effects to tests undertaken immediately after donning the orthosis.

537

## 538 **CONCLUSIONS**

539 The aims of this pilot RCT were partly achieved. Although the intended number and  
540 chronicity distribution of athletes was not reached, this may be addressed in the future by  
541 employing more focused recruitment drives (for example with Gaelic Football), extending the  
542 recruitment period and focusing upon athletes with chronic pain. The criteria of an attrition  
543 rate < 20% was achieved. The protocol itself was feasible, and blinding of the investigator  
544 was effective, but the use of co-investigators would be more time effective and essential for  
545 facilitating better recruitment.

546 The effect sizes and recruitment/dropout rates suggest that the intervention holds promise as a  
547 tool to support the multi-modality approach to pelvic / groin injury management. Based upon  
548 these findings and the actions proposed to address recruitment, a future definitively powered  
549 RCT appears feasible and is indicated.

550

## References

- 551  
552  
553 1. Glasgow P, Webb M, McNicholl C. *Report of Gaelic groin think tank: prevention and*  
554 *management of chronic groin pain in Gaelic Football*. Newtownabbey: Sports  
555 Institute Northern Ireland in association with Ulster Council GAA; 2011.
- 556 2. Tyler TFS, H.J.; Gerhardt, M.B.; Nicholas, S.J. Groin injuries in sports medicine.  
557 *Sports Health*. 2010;2(3):231-236.
- 558 3. Thorborg K, Rathleff MS, Petersen P, Branci S, Hölmich P. Prevalence and severity  
559 of hip and groin pain in sub-elite male football: a cross-sectional cohort study of 695  
560 players. *Scand J Med Sci Spor*. 2017;27(1):107-114.
- 561 4. Ficek K, Rzepka R, Orawczyk T, Zotnierzcyk Z. Groin pain in athletes - clinical  
562 experience. *J Hum Kinet*. 2008;19:141-148.
- 563 5. Ekçi B, Beyzadeoglu T. Groin pain in athletes - sports hernia and osteitis pubis. In:  
564 Canonico S, ed. *Inguinal Hernia [online]*. Vol 10.5772/58621 Intech. 111-123  
565 Available: [http://www.intechopen.com/books/export/citation/EndNote/inguinal-](http://www.intechopen.com/books/export/citation/EndNote/inguinal-hernia/groin-pain-in-athletes-sports-hernia-and-osteitis-pubis)  
566 [hernia/groin-pain-in-athletes-sports-hernia-and-osteitis-pubis](http://www.intechopen.com/books/export/citation/EndNote/inguinal-hernia/groin-pain-in-athletes-sports-hernia-and-osteitis-pubis) [Accessed 6/2/2015];  
567 2014.
- 568 6. Serner A, van Eijck CH, Beumer BR, Hölmich P, Weir A, de Vos R-J. Study quality  
569 on groin injury management remains low: a systematic review on treatment of groin  
570 pain in athletes. *Br J Sports Med*. 2015;49:818.  
571 <http://bjsm.bmj.com/content/49/12/813.full>. Accessed 12/2/ 2015.
- 572 7. Arumugam A, Milosavljevic S, Woodley S, Sole G. Effects of external pelvic  
573 compression on form closure, force closure, and neuromotor control of the  
574 lumbopelvic spine – a systematic review. *Manual Ther*. 2012;17(4):275-284.
- 575 8. Mens J, Inklaar H, Koes BW, Stam HJ. A new view on adduction-related groin pain.  
576 *Clin J Sport Med*. 2006a;16(1):15-19.

- 577 9. Sawle L, Freeman J, Marsden J, Matthews MJ. Exploring the effect of pelvic belt  
578 configurations upon athletic lumbopelvic pain. *Prosthetics and Orthotics*  
579 *International*. 2013;37(2):124-131.
- 580 10. Sawle L, Freeman J, Marsden J. The use of a dynamic elastomeric fabric orthosis  
581 (DEFO) in supporting the management of athletic pelvic and groin injury. *J Sports*  
582 *Rehabil*. 2016;25:101-110.
- 583 11. Beliard S, Chauveau M, Moscatiello T, Cros F, Ecarnot F, Becker F. Compression  
584 garments and exercise: no influence of pressure applied. *J Sports Sci Med*.  
585 2015;14(1):75-83.
- 586 12. Hill J, Howatson G, van Someren K, Leeder J, Pedlar C. Compression garments and  
587 recovery from exercise-induced muscle damage: a meta-analysis. *Br J Sports Med*.  
588 2013.
- 589 13. McKim KR, Taunton JE. The Effectiveness of compression shorts in the treatment of  
590 athletes with osteitis pubis. *NZ J Sport Med*. 2001;29:70-73.
- 591 14. Chaudhari AMW, Jamison ST, McNally MP, Pan X, Schmitt LC. Hip adductor  
592 activations during run-to-cut manoeuvres in compression shorts: implications for  
593 return to sport after groin injury. *J Sport Sci*. 2014;32(14):1333-1340.
- 594 15. Doan BK, Won YH, Newton RU, et al. Evaluation of a lower-body compression  
595 garment. *J Sport Sci*. 2003;21:601-610.
- 596 16. Kraemer WJ, Bush JA, Newton RU, et al. Influence of a compression garment on  
597 repetitive power output before and after different types of muscle fatigue. *Sports Med,*  
598 *Training and Rehab*. 1998;8(2):163-184.
- 599 17. Michael JS, Dogramaci SN, Steel KA, Graham KS. What is the effect of compression  
600 garments on a balance task in female athletes? *Gait & Posture*. 2014;39(2):804-809.

- 601 18. Arain M, Campbell MJ, Cooper CL, Lancaster GA. What is a pilot or feasibility  
602 study? A review of current practice and editorial policy. *BMC Medical Research*  
603 *Methodology*. 2010;10: 67:67. <http://www.biomedcentral.com/1471-2288/10/67>.  
604 Accessed 20/1/2013.
- 605 19. Thabane L, Ma J, Chu R, et al. A tutorial on pilot studies: the what, why and how.  
606 *BMC Medical Research Methodology*. 2010;10(1):1-10.
- 607 20. Orchard JW. Men at higher risk of groin injuries in elite team sports: a systematic  
608 review. *Br J Sports Med*. 2015;49(12):798-802.
- 609 21. Stuber KJ. Specificity, sensitivity, and predictive values of clinical tests of the  
610 sacroiliac joint: A systematic review of the literature. *The Journal of the Canadian*  
611 *Chiropractic Association*. 2007;51(1):30-44.
- 612 22. Vleeming A, Albert HB, Ostgaard HC, Sturesson B, Stuge B. European guidelines for  
613 the diagnosis and treatment of pelvic girdle pain. *European Spine Journal*.  
614 2008;17(6):794-819.
- 615 23. Hagglund M, Walden M, Ekstrand J. Injuries among male and female elite  
616 footballers. *Scand J Med Sci Spor*. 2009;19(6):819-827.
- 617 24. Elliott SA, Brown JSL. What are we doing to wait list controls? *Behaviour Research*  
618 *& Therapy*. 2002;40(9):1047-1052.
- 619 25. Schulz KF, Grimes DA. Blinding in randomised trials: hiding who got what. *The*  
620 *Lancet*. 2002;359(9307):696-700.
- 621 26. Dempster M. *A Research Guide for Health and Clinical Psychology*. Palgrave  
622 Macmillan; 2011.
- 623 27. Schulz KF, Altman DG, Moher D. CONSORT 2010 Statement: updated guidelines  
624 for reporting parallel group randomised trials. *BMC Medicine*. 2010;8(18):1-9.


- 625 28. Julious SA. Sample size of 12 per group rule of thumb for a pilot study.  
626 *Pharmaceutical Statistics*. 2005;4:287-291.
- 627 29. Wood A, White I, Thompson S. Are missing outcome data adequately handled? A  
628 review of published randomized controlled trials in major medical journals. *Clin*  
629 *Trials*. 2004;1:368 - 376.
- 630 30. Kottink AIR, Oostendorp LJM, Buurke JH, Nene AV, Hermens HJ, Ijzerman MJ. The  
631 Orthotic Effect of Functional Electrical Stimulation on the Improvement of Walking  
632 in Stroke Patients with a Dropped Foot: A Systematic Review. *Artif Organs*.  
633 2004;28(6):577-586.
- 634 31. Nevin F, Delahunt E. Adductor squeeze test values and hip joint range of motion in  
635 Gaelic football athletes with longstanding groin pain. *J Sci Med Sport*.  
636 2014;17(2):155-159.
- 637 32. Malliaras P, Hogan A, Nawrocki A, Crossley K, Schache A. Hip flexibility and  
638 strength measures: reliability and association with athletic groin pain. *Br J Sports*  
639 *Med*. 2009;43(10):739-744.
- 640 33. Hanna CM, Fulcher ML, Raina Elley C, Moyes SA. Normative values of hip strength  
641 in adult male association football players assessed by handheld dynamometry. *J Sci*  
642 *Med Sport*. 2010;13:299-303.
- 643 34. Hölmich P, Hölmich LR, Berg AM. Clinical examination of athletes with groin pain:  
644 an intraobserver and interobserver reliability study. *Br J Sports Med*. 2004;38:446-  
645 451.
- 646 35. Mens JMA, Vleeming A, Schniders CJ, Stam HJ, Ginai AZ. The active straight leg  
647 raise test and mobility of the pelvic joints. *European Spine Journal*. 1999;8:468-473.

- 648 36. Palsson TS, Hirata RP, Graven-Nielsen T. Experimental pelvic pain impairs the  
649 performance during the active straight leg raise test and causes excessive muscle  
650 stabilization. *Clin J Pain*. 2015;31(7):642-651.
- 651 37. Jansen JA, Mens JM, Backx FJ, Stam HJ. Changes in abdominal muscle thickness  
652 measured by ultrasound are not associated with recovery in athletes with longstanding  
653 groin pain associated with resisted hip adduction. *Journal of Orthopaedic and Sports  
654 Physical Therapy*. 2009;39(10):724-732.
- 655 38. Markovic G, Dizdar D, Jukic I, Cardinale M. Reliability and factorial validity of squat  
656 and countermovement jump tests. *J Strength Cond Res*. 2004;18(3):551-555.
- 657 39. Almuzaini KS, Fleck SJ. Modification of the standing long jump test enhances ability  
658 to predict anaerobic performance. *J Strength Cond Res*. 2008;22(4):1265-1272.
- 659 40. Riemann B, Caggiano NA, Lephart SM. Examination of a clinical method of  
660 assessing postural control during a functional performance task. *J Sports Rehabil*.  
661 1999;8:171-183.
- 662 41. Sawle L, Freeman J, Marsden J. Intra-rater reliability of the multiple single-leg hop-  
663 stabilization test and relationships with age, leg dominance and training. *Int J Sports  
664 Phys Ther*. 2017;12(2):190-198.
- 665 42. Velotta J, Weyer J, Ramirez A, Winstead J, Bahamonde R. Relationship between leg  
666 dominance tests and type of task. *Portugese Journal of Sports Sciences*.  
667 2011;11(2):1035-1038.
- 668 43. Freeman JV, Julious SA. The analysis of categorical data. *Scope*. 2007;16(1):18-21.
- 669 44. Lancaster GA, Dodd S, Williamson PR. Design and analysis of pilot studies:  
670 recommendations for good practice. *J Eval Clin Pract*. 2004;10(2):307-312.
- 671 45. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Hillsdale:  
672 Lawrence Erlbaum Associates; 1988.

- 673 46. Gupta SK. Intention-to-treat concept: a review. *Perspect Clin Res*. 2011;2(3):109-112.
- 674 47. Streiner D, Geddes J. Intention to treat analysis in clinical trials when there are  
675 missing data. *Evidence Based Mental Health*. 2001;4(3):70-71.
- 676 48. Bhakta BB. Management of spasticity in stroke. *Brit Med Bull*. 2000;56(2):476-485.
- 677 49. Beales DJ, O'Sullivan PB, Briffa NK. The effects of manual pelvic compression on  
678 trunk motor control during an active straight leg raise in chronic pelvic girdle pain  
679 subjects. *Man Ther*. 2010;15(2):190-199.
- 680 50. Bernhardt T, Anderson GS. Influence of moderate prophylactic compression on sport  
681 performance. *J Strength Cond Res*. 2005;19(2):292-297.
- 682 51. Born DP, Holmberg HC, Goernert F, Sperlich B. A novel compression garment with  
683 adhesive silicone stripes improves repeated sprint performance - a multi-experimental  
684 approach on the underlying mechanisms. *BMC Sports Sci Med Rehabil*. 2014;6:21.
- 685 52. Mostert-Wentzel K, Swart JJ, Masenyetse LJ, et al. Effect of kinesio taping on  
686 explosive muscle power of gluteus maximus of male athletes: original research. *South  
687 African Journal of Sports Medicine*. 2012;24(3):75-80.
- 688 53. Hertel J, Miller JS, Denegar CR. Intratester and intertester reliability during the star  
689 excursion balance tests. *J Sport Rehabil*. 2000;9:104-116.
- 690 54. Kirk RE. *Experimental Design: Procedures for the Behavioral Sciences*. 2nd ed.  
691 Pacific Grove: Brooks/Cole; 1982.
- 692 55. Jung HS, Jeon HS, Oh DW, Kwon OY. Effect of the pelvic compression belt on the  
693 hip extensor activation patterns of sacroiliac joint pain patients during one-leg  
694 standing: a pilot study. *Man Ther*. 2013;18(2):143-148.
- 695 56. Whittaker JL, Small C, Maffey L, Emery CA. Risk factors for groin injury in sport: an  
696 updated systematic review. *Br J Sports Med*. 2015;49:803-809.

- 697 57. Venckunas T, Trinkunas E, Kamandulis S, Poderys J, Grunovas A, Brazaitis M.  
698 Effect of lower body compression garments on hemodynamics in response to running  
699 session. *Scientific World Journal*. 2014;353040:353040.  
700 <http://www.hindawi.com/journals/tswj/2014/353040/>. Accessed 15/2/2015.
- 701 58. Racinais S, Oksa J. Temperature and neuromuscular function. *Scand J Med Sci Spor*.  
702 2010;20(s3):1-18.
- 703
- 704






705

706 **Figure 1 Athlete pathway through the study**

707



**Figure 2 The CONSORT diagram showing the flow of participants through the study**

708

709

	<b>Waiting-List Control Group</b> <b>(n = 9)</b>	<b>Intervention</b> <b>Group</b> <b>(n = 7)</b>
Gender	Male = 6	Male = 7
Leg Dominance	Right = 8	Right = 7
Mean Age in years +/-SD (range)	30.7 +/- 9.3 (22-48)	26 +/- 5.3 (23-36)
Mean Height in cm +/- SD (range)	179 +/- 6.2 (167-190.5)	180 +/- 8 (164.8-186.5)
Mean Weight in kg +/- SD (range)	73.2 +/- 15 (56.4-93.4)	80.5 +/- 7.8 (66.2-88.7)

710

711 **Table 1 Athlete demographics**

712

	Week 2		Week 4		Week 6	
<b>Outcome Measure</b>	<b>Compression Orthosis Effect Size (d)</b>	<b>Sport Shorts Effect Size (d)</b>	<b>Compression Orthosis Effect Size (d)</b>	<b>Sport Shorts Effect Size (d)</b>	<b>Compression Orthosis Effect Size (d)</b>	<b>Sport Shorts Effect Size (d)</b>
<b>Dominant Leg ASLR NRS Score</b>	0.4	0.2	0.7 <sup>^</sup>	0.4	0.6 <sup>^</sup>	0.5 <sup>^</sup>
<b>Dominant Leg ASLR Score</b>	0.7 <sup>^</sup>	0.3	0.8 *	0.4	1.1*	0.7 <sup>^</sup>
<b>Non-Dominant Leg ASLR NRS Score</b>	0.6 <sup>^</sup>	0.8*	0.8*	0.6 <sup>^</sup>	0.9*	0.8*
<b>Non-Dominant Leg ASLR Score</b>	0.7 <sup>^</sup>	0.6 <sup>^</sup>	0.2	0.6 <sup>^</sup>	1.1*	0.8*
<b>Squeeze Test Force (N)</b>	- 0.1	0.0	0.8*	0.6 <sup>^</sup>	0.8*	-0.5
<b>Broad Jump Distance (cm)</b>	0.1	-0.1	0.2	0.1	0.2	0.1
<b>Dominant Leg MSLHST Score</b>	- 0.1	0.1	0.2	0.5 <sup>^</sup>	0.1	-0.4
<b>Non-Dominant Leg MSLHST Score</b>	0.1	-0.3	0.8*	0.7 <sup>^</sup>	0.3	0.0

713

714 **Table 2 The effect sizes (d) for each outcome measure at each stage of the study.**

715 NRS refers to the numerical rating scale; MSLHST refers to the functional balance test

716 \*Denotes a large effect size; <sup>^</sup> signifies a moderate effect size

717

	Intervention Group Mean Difference and 95% Confidence Intervals						Waiting-List Control Group Mean Difference and 95% Confidence Intervals		
	DEFO			Sport Shorts			Sport Shorts		
Measures	Week Two	Week Four	Week Six	Week Two	Week Four	Week Six	Week Two	Week Four	Week Six
<b>Dominant Leg ASLR NRS Score</b>	<b>1.1</b> -1.4 3.6	<b>1.3</b> -0.9 3.4	<b>1.2</b> -1.0 3.3	<b>0.8</b> -1.6 3.1	<b>1.0</b> -1.3 3.2	<b>1.1</b> -1.1 3.3	<b>0.4</b> -0.8 1.7	<b>0.3</b> -1.0 1.5	<b>0.2</b> -1.4 1.8
<b>Dominant Leg ASLR Mens Score</b>	<b>0.6</b> -0.4 1.7	<b>0.6</b> -0.4 1.7	<b>0.8</b> -0.1 1.7	<b>0.4</b> -0.7 1.4	<b>0.4</b> -0.8 1.5	<b>0.5</b> -0.4 1.4	<b>0.1</b> -0.5 0.7	<b>0.0</b> -0.8 0.7	<b>0.0</b> -0.6 0.7
<b>Non-Dominant Leg ASLR NRS Score</b>	<b>1.2</b> -1.2 3.5	<b>1.4</b> -0.8 3.6	<b>1.3</b> -1.0 3.5	<b>1.4</b> -0.8 3.6	<b>1.1</b> -1.1 3.3	<b>1.1</b> -1.0 3.3	<b>0.2</b> -1.0 1.3	<b>0.2</b> -0.8 1.3	<b>-0.3</b> -1.9 1.4
<b>Non-Dominant Leg ASLR Mens Score</b>	<b>0.5</b> -0.6 1.6	<b>0.1</b> -1.4 1.6	<b>0.7</b> -0.3 1.7	<b>0.4</b> -0.8 1.6	<b>0.4</b> -0.7 1.5	<b>0.4</b> -0.7 1.5	<b>-0.2</b> -0.9 0.6	<b>-0.1</b> -0.8 0.6	<b>-0.2</b> -1.1 0.6
<b>Squeeze Test Force</b>	<b>-5.6</b> -72.5 61.4	<b>43.6</b> -30.3 117.4	<b>86.8</b> 21.6 152.0	<b>-1.3</b> -73.1 70.5	<b>27</b> -47.9 101.8	<b>-0.8</b> -52.0 50.4	<b>0.7</b> -70.1 71.5	<b>-9.7</b> -78.1 58.7	<b>32.4</b> -42.8 107.6

<b>Broad Jump Distance</b>	<b>3.6</b> -36.1 43.2	<b>9.9</b> -29.4 49.1	<b>11.8</b> -26.1 49.7	<b>-4.4</b> -43.7 34.8	<b>5.8</b> -38.9 50.4	<b>9.2</b> -33.9 52.2	<b>0.0</b> -39.0 39.0	<b>2.3</b> -39.2 43.8	<b>4.0</b> -37.8 45.9
<b>Dominant Leg MSLHST Error Score</b>	<b>-1.5</b> -20.0 17.0	<b>0.9</b> -16.9 18.7	<b>10.9</b> -6.9 28.7	<b>2.5</b> -14.2 19.2	<b>6.2</b> -10.3 22.7	<b>3.1</b> -13.7 19.8	<b>1.2</b> -20.6 23.1	<b>-1.9</b> -18.8 25.4	<b>9.1</b> -11.2 29.3
<b>Non- Dominant Leg MSLHST Error Score</b>	<b>5.3</b> -10.1 20.7	<b>10.7</b> -4.3 25.7	<b>10.6</b> -4.3 25.4	<b>-2.3</b> -20.6 16.0	<b>9.0</b> -11.2 29.2	<b>6.6</b> -14.9 28.1	<b>3.3</b> -20.6 16.7	<b>-3.8</b> -25.2 17.5	<b>6.5</b> -11.7 24.7

718

719 **Table 3. The mean difference (in bold) and 95% confidence intervals for the mean difference, from baseline to assessment week two,**  
720 **four and six for each outcome measure and for each condition**

721

