



UNIVERSITY OF
PLYMOUTH

PEARL

QOL-05. E-SURVEY OF CURRENT INTERNATIONAL PHYSIOTHERAPY PRACTICE FOR CHILDREN WITH ATAXIA FOLLOWING SURGICAL RESECTION OF POSTERIOR FOSSA TUMOUR

Hartley, H; Carter, B; Bunn, LM; Kumar, R; Cassidy, E; Lane, S; Pizer, B

Published in:

June 2018 *Neuro-Oncology* 20(suppl_2):i158-i158 DOI: 10.1093/neuonc/noy059.587

DOI:

[10.1093/neuonc/noy059.587](https://doi.org/10.1093/neuonc/noy059.587)

Publication date:

2018

Link:

[Link to publication in PEARL](#)

Citation for published version (APA):

Hartley, H., Carter, B., Bunn, LM., Kumar, R., Cassidy, E., Lane, S., & Pizer, B. (2018). QOL-05. E-SURVEY OF CURRENT INTERNATIONAL PHYSIOTHERAPY PRACTICE FOR CHILDREN WITH ATAXIA FOLLOWING SURGICAL RESECTION OF POSTERIOR FOSSA TUMOUR. *June 2018 Neuro-Oncology* 20(suppl_2):i158-i158 DOI: 10.1093/neuonc/noy059.587, 0(0). <https://doi.org/10.1093/neuonc/noy059.587>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Download date: 15. Feb. 2025



The ASPECT Study- E-survey of current international physiotherapy practice for children with ataxia following surgical resection of posterior fossa tumour

H Hartley¹, B Carter^{1,2}, L Bunn³, R Kumar¹, E Cassidy⁴, S Lane⁵, B Pizer¹.

¹Alder Hey Children's NHS Foundation Trust, ²Edge Hill University, ³University of Plymouth, ⁴Freelance Academic, ⁵University of Liverpool.

Background

Access to neuro-rehabilitation is recognized as crucial in paediatric neuro-oncology¹ with physiotherapy integral to this; yet there is no consensus as to the type, intensity or timing of interventions. In particular, there is a gap in the literature regarding physiotherapy intervention for children with ataxia following surgical resection of posterior fossa tumour. An e-survey was carried out to provide information regarding current international practice and guide areas of further research.

Objective

To determine current international practice regarding physiotherapy input for children with posterior fossa tumours.

Methods


An e-survey was developed covering the following domains;

- respondent demographics
- physiotherapy treatment and intervention
- virtual training
- intensity/timing of treatment
- outcomes of physiotherapy treatment.

The e-survey was piloted and refined, then distributed to physiotherapists via 6 key groups; Paediatric Oncology Physiotherapy Network (POPs), Association of Paediatric Physiotherapists (APCP), European Paediatric Neurology Society (EPNS), International Society of Paediatric Oncology (SIOP), Posterior Fossa society (PFS) and Pediatric Oncology Special Interest Group (American Physical Therapy Society).

Results

- 96 physiotherapists responded (UK n=53, rest of Europe n=23, USA/Canada n=10, Aus/NZ n=10).
- 41% of respondents had over 10 years experience in treating children with brain tumours.
- Respondents indicated they used a range of therapy interventions. The most commonly used were balance exercises (n=73), gait re-education (n=71), proximal control activities (n=70), and task specific training (n=65).
- Frequently used adjuncts to treatment were mobility aids and orthotics.
- Three themes were identified regarding challenges to physiotherapy input; condition specific factors (direct medical and treatment related), child and family factors (psychosocial and expectations/engagement) and physiotherapy delivery factors (lack of evidence and resources).



Conclusions

This e-survey provides an initial scoping review of international physiotherapy practice. It demonstrates the wide range of intervention types used and highlights the lack of clinical evidence in this area. The results raise the need for further research in this field to help with the development of physiotherapy guidelines in children with posterior fossa tumours.

¹ National Institute for Health and Clinical Excellence (2014) Cancer Services for children and young people. Quality Standard QS55.