



PEARL

Patient-to-Patient Interactions During the Pain Management Programme: The Role of Humor and Venting in Building a Socially Supportive Community

Anil, Krithika

Published in:

Frontiers in Pain Research

DOI:

[10.3389/fpain.2022.875720](https://doi.org/10.3389/fpain.2022.875720)

Publication date:

2022

Link:

[Link to publication in PEARL](#)

Citation for published version (APA):

Anil, K. (2022). Patient-to-Patient Interactions During the Pain Management Programme: The Role of Humor and Venting in Building a Socially Supportive Community. *Frontiers in Pain Research*, 3(0). <https://doi.org/10.3389/fpain.2022.875720>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Author's Proof

Before checking your proof, please see the instructions below.

- Carefully read the entire proof and mark all corrections in the appropriate place, using the Adobe Reader commenting tools ([Adobe Help](#)).
- Provide your corrections in a single PDF file or post your comments in the Production Forum making sure to reference the relevant query/line number. Upload or post all your corrections directly in the Production Forum to avoid any comments being missed.
- We do not accept corrections in the form of edited manuscripts nor via email.
- Do not provide scanned, handwritten corrections.
- Before you submit your corrections, please make sure that you have checked your proof carefully as once you approve it, you won't be able to make any further corrections.
- To ensure the timely publication of your article, please submit the corrections within 48 hours. After submitting, do not email or query asking for confirmation of receipt.

Do you need help? Visit our [Production Help Center](#) for more information. If you can't find an answer to your question, contact your Production team directly by posting in the Production Forum.

Quick Check-List

- Author names** - Complete, accurate and consistent with your previous publications.
- Affiliations** - Complete and accurate. Follow this style when applicable: Department, Institute, University, City, Country.
- Tables** - Make sure our formatting style did not change the meaning/alignment of your Tables.
- Figures** - Make sure we are using the latest versions.
- Funding and Acknowledgments** - List all relevant funders and acknowledgments.
- Conflict of Interest** - Ensure any relevant conflicts are declared.
- Supplementary files** - Ensure the latest files are published and that no line numbers and tracked changes are visible.
Also, the supplementary files should be cited in the article body text.
- Queries** - Reply to all typesetters queries below.
- Content** - Read all content carefully and ensure any necessary corrections are made.



Patient-to-Patient Interactions During the Pain Management Programme: The Role of Humor and Venting in Building a Socially Supportive Community

Katherine A. Finlay^{1*}, Adam Madhani¹, Krithika Anil² and Sue M. Peacock³

¹ School of Psychology and Clinical Language Sciences, University of Reading, Reading, United Kingdom, ² Peninsula Allied Health Centre, School of Health Professions, University of Plymouth, Plymouth, United Kingdom, ³ Pain in the Mind, Independent Practice, The Saxon Clinic, Milton Keynes, United Kingdom

OPEN ACCESS

Edited by:

Michele Trimboli,
Azienda Ospedaliero-Universitaria
Mater Domini, Italy

Reviewed by:

Margaret Dunham,
Edinburgh Napier University,
United Kingdom
Letizia Troisi,
University of Magna Graecia, Italy

*Correspondence:

Katherine A. Finlay
katherine.finlay@reading.ac.uk

Specialty section:

This article was submitted to
Pain Mechanisms,
a section of the journal
Frontiers in Pain Research

Received: 14 February 2022

Accepted: 21 March 2022

Published: xx March 2022

Citation:

Finlay KA, Madhani A, Anil K and
Peacock SM (2022) Patient-to-Patient
Interactions During the Pain
Management Programme: The Role of
Humor and Venting in Building a
Socially Supportive Community.
Front. Pain Res. 3:875720.
doi: 10.3389/fpain.2022.875720

Objectives: Social support is most positively perceived when there is an optimal match between a patient's need for communication and the purpose of their interaction. Maladaptive communication patterns may inhibit social bonding or mutual support, negatively impacting clinical outcomes. This study aimed to identify how people with chronic pain naturalistically converse together about their pain in the context of a Pain Management Programme (PMP).

Methods: Seven participants (4 females; 3 males) with ongoing chronic pain who were attending a PMP in a regional hospital in the United Kingdom were audio/video recorded during breaks in their PMP. Interactions were transcribed using Jeffersonian Transcription and analyzed using Conversation Analysis.

Results: Two conversational mechanisms were identified: (1) *Conversational humor*; and (2) *A venting cycle*. Participants used their pain-related experiences construct a motive for a joke, then proceeded to deliver the joke, which initiated a joke return from observers. The sequence was completed by a collaborative punchline. In the venting cycle, an initial complaint was escalated by the sharing of comparable experiences, after which the vent was concluded through a joke punchline, acting as a pivot to move the conversation forwards, terminating the venting.

Conclusions: Humorous interpersonal interactions about chronic pain provided a forum for social support-building within the PMP. Humor was affiliative and built social collaboration, helping individuals to together make sense of their pain in a prosocial atmosphere, approaching pain-related experiences with levity. Patient-to-patient interactions within the PMP were strongly prosocial and inclusive, potentially facilitating enhanced PMP clinical outcomes through collaboration.

Keywords: chronic pain, peer, social support, interpersonal, interaction, conversation analysis, qualitative

INTRODUCTION

The prevalence of chronic pain (CP), persisting for more than 3 months, is rising globally (1). Recent pooled estimates suggest that 34% of adults in the United Kingdom currently experience chronic pain, with chronic widespread pain affecting 5.5 million (12%) of the British population (2). The impact of CP is multifactorial, and aside from its physical and psychological impact, it places significant social burden on People Living with Chronic Pain (PLwCP) and those around them. Factors such as pain-related loss of employment (3) and withdrawal from social engagement (4) are often linked with a significant deterioration in quality of life for PLwCP and their families (5). As a result, the social environment surrounding PLwCP has been widely investigated: where lack of social support is strongly associated with the genesis, maintenance, and worsening of pain symptoms for PLwCP (6), by contrast, positive social relationships and support from family and friends builds work ability, psychological well-being and quality of life (7, 8). The salience of social support when living with CP is therefore clearly evident and it is important that PLwCP are supported to maintain and/or build positive social relationships and social support structures.

Yet recent research has suggested that not all (positive) social support is equal. In line with the optimal matching hypothesis (9), when social support is optimally matched to a person's needs, then their ability to cope with pain is increased (7, 8). Indeed, CP patients report increased satisfaction and lower perceived stress when expressing their feelings to PLwCP who are experiencing similar health challenges (7, 10). However, when there is a mismatch between an individual's health status and the support which is received, this support can be experienced as "toxic positivity" or unsolicited advice (7). This may mean that individuals feel obligated to generate false positivity, expressed and experienced by people with and without CP as inauthentic, reducing individuals' desires to offer or accept social support in the future.

Though social support "matching" undoubtedly confers some benefits, it is possible that the associated outcomes may not be universally positive. Research has demonstrated that when individuals experiencing similar situations express their perspectives collectively, negative emotional contagion may occur (11). This phenomenon reflects the transfer of negative moods and emotions within a group (12), and the automatic synchronization and convergence of such emotions (13). By extension, in the context of chronic pain, this negative exchange spiral of emotions may increase perceived pain symptoms and lead to an overall decline in well-being. Though emotional contagion specifically within chronic pain settings has not yet been investigated, there is undoubtedly a need to better understand the formulation and presentation of person-to-person interactions in PLwCP. Chronic pain management interventions are typically undertaken in group contexts (14) and it is possible that group interactions within these settings may be jeopardizing rather than enhancing outcomes of pain management programmes.

An emerging method to understand emotion and pain experiences is a discursive approach (15). Discursive approaches move beyond pain self-report measures, to look at how "actions in talk" can better depict pain experiences (16). The analytic method of conversation analysis (CA) is a strong discursive approach which gathers information about normative patterns of speech in conversation and it analyses the extent to which deviant cases occur (17). CA is a novel way of understanding medical and clinical interactions (18), and has been widely used in clinical settings to understand communication between patients and clinicians (19–21).

To date, no studies have used CA to assess patient-to-patient interactions; instead, CA research typically focuses on clinician-patient interactions, addressing issues such as resistance (22) and advice giving (23) CA is often exploited to explore conversations on sensitive topics such as weight management (24) and cancer (25), but has not yet been extended to use in chronic pain. Consequently, the application of CA in a chronic pain setting may prove effective in understanding the conversation sequences and discursive organization presenting in interactions between PLwCP. CA may facilitate exploration of the conversational interactions involved in seeking social support from individuals also living with CP within a pain management group setting. This study therefore aimed to apply CA to patient-to-patient interactions in chronic pain management programmes, to investigate the conversational mechanisms involved in seeking mutual support between PLwCP.

METHOD

Design

A qualitative observational study design was employed, using Conversational Analysis in the context of a 6 week Pain Management Programme.

Participants and Recruitment

This study was approved by the University of Buckingham School of Psychology and Well being Ethics Committee and was granted full NHS ethical approval (REC reference 13/WM/0214). Six weeks before the PMP began, all attendees at three prospective Pain Management Programmes were invited to take part in this study by letter. Inclusion criteria were participants above the age of 18 years, who had experienced chronic pain for at least 3 months, and had attended two or more seminars at the Pain Management Unit, to gain eligibility for PMP attendance. Chronic pain presentation for each patient was confirmed prior to recruitment for this study by objective clinical and subjective pain assessment measures, undertaken during routine clinical care by the specialist Multidisciplinary Team in the Pain Management Unit (comprising Doctors, Pain Consultants, Physiotherapists, Occupational Therapists, Nurses and Psychologists) before referral to the Pain Management Programme.

Preliminary consent to participation was obtained from 19 patients (8 males and 11 females) across three PMPs. Although consent was obtained from 19 volunteers, in only one out of the three PMP groups was consent to filming/audio recording given

by all PMP group members. Therefore, to ensure anonymity of non-consenting PMP members, data was only recorded from one PMP ($N = 7$; 4 females; 3 males), in which participant ages ranged between 33 and 77 years. All participants were Caucasian and living with chronic, non-malignant, musculoskeletal pain in the lower back (five participants), legs (one participant) or neck (one participant).

Pain Management Programme Overview

The PMP was a 6 week programme, comprising two 4-h sessions per week, for adults with non-malignant chronic pain. The PMP was delivered face-to-face in a Secondary Care setting (Milton Keynes University Hospital) in the United Kingdom by a multidisciplinary team of pain management specialists. The PMP was undertaken in accordance with the Recommended Guidelines for Pain Management Programmes for Adults (26). The content of the PMP included pain psychoeducation and teaching on biological mechanisms of pain, goal setting, activity pacing, stress management, relaxation techniques, re-engaging with physical activity, cognitive behavioral approaches, self-monitoring, sleep hygiene, the benefits of social support, managing pain medication and dealing with pain flare-ups (26).

Procedure

Participants were given a short presentation at the start of the first pain management programme session, highlighting the study's aim and design and were encouraged to ask questions about the research. Participants who volunteered to take part were provided with a consent form and written, and informed consent was obtained. Participants were informed about their right to withdraw at any point, without facing negative consequences or affecting their involvement in the PMP. It was explained that as the study was investigating naturalistic conversation in pain settings, the recording equipment (video and audio) would be set up only in the coffee room and recordings would take place during scheduled breaks between PMP teaching sessions. This was to ensure that specific personal/clinical information or Healthcare Professional-led teaching was not recorded, and patient-to-patient interactions were prioritized, in accordance with the aims of the study. Recordings could only be made when fully consenting participants were together and conversing freely without third party involvement during their PMP breaks. In total, data were viable to be collected from four 20 min coffee break sessions across 8 weeks, which resulted in 78 min of conversation of recorded data. All data gathered were stored on a password-protected laptop and external hard drive. At the end of the study, the participants were fully debriefed and signposted to organizations offering further support with managing chronic pain.

Data Preparation and Analysis

Recordings were transcribed using Jeffersonian transcription techniques (27) and were analyzed using CA. Jeffersonian transcription method differs from verbatim transcription as it highlights talk in greater detail to understand vocal nuances, which are then included in the analysis (See **Table 1** for transcription symbols). Jefferson (27) organizes in-talk

TABLE 1 | Transcription symbols adapted from Jefferson (27).

Timing of utterances		
Equals sign	=	No interval between adjacent utterances
Time in single brackets	(0.5)	Length in tenths of seconds of pause in the talk
Full stop in single brackets	(.)	Pause that is less than one tenth of a second
Left hand square bracket	[Beginning point of an overlap in utterance
Right hand square bracket]	Ending of an overlap in utterance
Speech delivery symbols		
Colon	:	Extension of a sound or syllable it follows
Full stop	.	Stopping fall in tone. Does not necessarily indicate end of a sentence
Comma	,	A continuing intonation
Question mark	?	Rising inflection, not necessarily a question
Exclamation mark	!	Animated tone, not necessarily an exclamation
Single dash	-	Halting or abrupt cut off to a word
Upward arrow	↑	Rise in pitch
Downward arrow	↓	Fall in pitch
Underline	<u>Emphasis</u>	Talk that is emphasized
Capital letters	LOUD	Talk that is louder than the surrounding
Degree sign	°quiet°	Talk that is quieter than the surrounding
Heh	Heh	Aspiration or laughter
'h' in brackets	(h)	Noticeable aspiration or laughter within speech
Full stop before 'h'	.h	Inbreath
h or series of hhh	h/hhh	Outbreath
Lesser than/greater than signs	> talk <	Talk delivered at greater speed
Greater than/lesser than signs	< talk >	Talk delivered at slower speed / elongated talk
Empty bracket	()	Inaudible transcript
Word or phrase in brackets	(talk)	Transcription in doubt
Words in double brackets	(laughs)	Description of sound

interaction transcription convention into five categories: (1) transcript layout, (2) temporal and sequential relationships, (3) speech delivery, which includes tone, pitch, tempo, and emphasis, (4) metacommentary and uncertain hearings and (5) representation of non-verbal speech such as laughing and crying. Data was transcribed in full for all conversational interactions.

All identifiable features such as names and places were anonymised during the transcription process. Transcription was carried out independently by the first and second author. Final transcription was agreed upon by both authors through discussion and clarification of discrepancies by repeated hearings of audio data, supplemented by video data if required. Once

transcription was finalized, all authors independently carried out an analysis to identify speech patterns within the data. Comparison of analysis patterns were discussed within the research group and commonalities and differences identified. The final analysis framework was therefore the result of both independent analyses and group discussion and was confirmed only when agreement was reached by all authors (28). The representativeness of the framework was further evaluated by presenting the final data to a CA specialist with significant experience of discursive psychology approaches. Through discussion, all discrepancies were resolved until consensus was reached amongst authors and the final coding framework of speech patterns was finalized.


Conversation Analysis: Theoretical Stance and Key Conversational Mechanisms

CA aims to examine the sequence organization of talk and the social actions that it can achieve (29). It is grounded in the ethnomethodology theoretical framework and explores the detailed organization of interactions in a naturalistic setting (25). The most basic mechanism explored in CA revolves around turn-taking (30). Like other forms of coordinated activities, conversations require turn-taking to ensure organization between participants (30). Therefore, the validity of CA results emerges from focusing on how each successive turn provides evidence for how the next speaker interprets and understands the speaker (31). Additionally, Bateson (32) argues that it is possible to frame a conversation as serious or playful by signaling to the other members the playful nature of the speech. When conversations are framed within a play frame (32) participants are aware that the ongoing conversation is defined as play and participants in the conversation should interpret it like this. Finally, backchannel responses and the display of positive politeness are common in everyday conversations. Positive politeness (33) is expressed by showing similarities amongst each other and expressing appreciation and understanding of others' experiences (34). Additionally, agreement and solidarity toward a speaker is also observed through repetition of speaker's thoughts and feelings (33). Backchanneled minimal responses such as "mhmm" and "yeah" can demonstrate a form of agreement between participants (35). These mechanisms observed in conversation all help understand and frame the sequence and dynamics between individuals while interacting with each other.

RESULTS

The results of this study demonstrated that two key conversational mechanisms were active when PLwCP were interacting with each other and exploring their mutual experiences of chronic pain (see **Figures 1, 2**):

- 1) *Conversational humor*: The use of humor in naturalistic patient-to-patient conversation to share and validate pain experiences.
- 2) *The Venting cycle*: Catharsis of negative emotions and difficult experiences through expressing personal concerns which are



received, expanded, and concluded by listeners in patient-to-patient interactions.

Conversational Humor

Conversational humor was employed by all participants when directly discussing their personal pain experiences and pain management. The humor followed a turn-taking sequence: participants set up a joke by providing a contextual "motive," which led into the initiation of a joke, triggering an interactive response (joke return) and the humor sequence was then closed by a punchline. Exemplars of conversational humor are demonstrated in the three extracts presented below (see **Tables 3–6**).

Extract 1: Conversational Humor

In Extract 1 (**Table 2**) a conversation is undertaken between three participants who joke about the amount of medication they need to take as individuals who experience chronic pain.

In this extract, Tyrone sets up the conversational joke by comparing the number of tablets his girlfriend takes to himself (Lines 1–3). The joke is presented when the patient says: "I'm taking 11 a day: y? now" (Line 6). The positioning of the utterance "hehe" in both lines 4 and 7 indicates the levity recognized within the conversation (Schenein, 1972), suggesting that the conversation is set within a play frame. Brenda then initiates the joke's return by suggesting they take even more medication than Steve (Line 9). Tyrone responds with "Eh come 'ere come 'ere" as the joke return. Comedic miming accompanied this conversational joke return, emphasizing the nature of the joke. Brenda then begins a new joke within the same context by referring to tablets as "sweeties" (Line 14). Tyrone returns the joke and links the two jokes by again saying, "Come 'ere come 'ere eh come 'ere." Finally, both Tyrone and Brenda, who have been the core participants in the joke and joke-return cycle in this extract, contribute together to the joke's punchline. To conclude the joking, Brenda describes how they have consumed enough medication to have a chemist inside them (Line 17 and 18), and Tyrone adds to this punchline by saying: "I could pick you up and rattle yeh" (Line 19).

From this extract, it is seen that rather than making a joke directed toward each other or teasing one another, the participants joke together about their own experiences and collectively share their experiences through humor. This cycle continues until the joke punchline. In this example, both individuals contribute to the punchline. The third participant in the conversation acted as a bystander who validated the play frame and collaboratively participated in the humorous nature of the conversation by uttering




TABLE 2 | Extract 1: Joking cycle.

Line number	Pseudonym	Conversation text	Conversation analysis
1	Tyrone:	Girlfriend's got ai ai uh ti tendonitis and arthritis(.	Set up of joke (Motive)
2		and I think she's taking seven tablets a day and then when I	
3		turned around to 'em and said o:h I GOT MORE TABLETS OFF UH	
4		OF A DOCTOR and she said (.) hehe	
5		>how many you taking now and I said<	Joke 1 presented
6		i'm taking 11 a da:y? now	
7	Steve:	Hehhh	Conversation continues and build up to return
8	Tyrone:	> 11 a day I'm taking < ↓ right	
9	Brenda:	16!	
10	Tyrone:	16! are you	Joke return
11	Brenda:	Aye (.)	
12	Tyrone:	[Eh come 'ere come 'ere]	Joke 2 presented
13	Brenda:	[I tell 'em when I when I come in and get my tablets] I say	
14		Can I come for my sweeties? now	Joke 2 return
15	Tyrone:	[Come 'ere come 'ere] eh come 'ere	
16	Steve:	[Hehhhh]	Minimal response
17	Brenda:	And I said to her >if you open me up you'll find< th	Joke punchline
18		chemist inside	
19	Steve:	I could pick you up and rattle yeh Heh	

verbal indicators of amusement: “hehe.” Even though Steve does not interact with the conversation in-depth, the minimal response of “hehe” acts as a positive indicator that he is paying positive attention to the speaker (36) and engaging in active listening and emotional expressions of collaborative humor.

Extract 2: Conversational Humor

The extract in Table 3 demonstrates a second exemplar which uses the same conversational humor speech pattern. In this extract, patients joke about trialing medications and medical marijuana. Four participants are present in this conversation. However, only two (Gemma and Steve) are engaging with each other, and the other two are bystanders who are listening to the conversation.

Gemma provides the joke’s motive by discussing a medical trial program that she is on through a University. Steve demonstrates empathetic engagement toward the situation in lines 213 and 216 as Gemma describes that she is on this program as no other treatment is working. Gemma then initiates the joke in describing a postman bringing the medication: “so my

postman arrives with my black bag” (Line 219). The minimal response by Brenda and Steve with the exhaled “hehh” utterance signifies the play frame and joke acknowledgment. In saying “You mean a suitcase?” (Line 223), Steve provides a return to Brenda’s joke. This joke adds to the empathetic response that was given to Gemma by Steve earlier in the conversation, demonstrating social support through the use of conversational humor. The joking cycle continues with Gemma initiating another joke as a response to the previous joke return: “Yeah ↑you meet a bloke around the back of an alley” (Line 223). Once again, the other two participants’ minimal responses indicate conversational engagement, and they participate in validating the collaborative humor within the conversation (Lines 225 and 226). Gemma continues the joke: “So yeah Yeah hehhhh I’d get stopped by the police.” (Line 227) and finally, both Gemma and Steve contribute to the punchline. Gemma says, “they’re prescription it’s alright hehh” referring to the police catching the medication, and Steve extends this punchline by adding “You know I’m smoking marijuana but eh given to me by my GP.” As the joke was steering toward illegal medication, Steve adds to the punchline by adding ironic clarification: in the hypothesized situation built within

TABLE 3 | Extract 2 Joking cycle.

Line number	Pseudonym	Conversation text	Conversation analysis
211	Gemma:	That's through (.) London it's through uh King's College in London that's through °so yeah I'm on trial for that°	Set up of joke (Motive)
212		London that's through °so yeah I'm on trial for that°	
213	Steve:	°Like you're being treated like a guinea pig for that so°	
214	Gemma:	Yeah h. but that was the step I had to take	Continuation of joke set up
215		[No other medicine was working]	
216	Steve:	[Yeah I'm just saying you had no other choice]	
217	Gemma:	And I was like well if it's going to make me a <little bit> better then I'll give it a go	
218		So my postman arrives with my black bag	Joke 1 presented
219		Hehh	Minimal responses
220	Brenda:	Hehhh	
221	Steve:	Hehhh	
222	Gemma:	It's my medication so I said as long as I-	Setup of return
223	Steve:	You mean a suitcase?	Joke return
224	Gemma:	Yeah ↑you meet a bloke around the back of an alley	Joke 2 presented
225	Steve:	Hehh	Minimal responses
226	Brenda:	Hehh	
227	Tyrone:	Yeah	
228	Gemma:	So yeah Yeah hehhhh I'd get stopped by the police.	Joke return
229	Steve:	Hehh	Minimal response
230	Gemma:	!They're prescription it's alright hehh	Joke punchline
231	Steve:	You know I'm smoking marijuana but eh hhh given to me by my GP hehh	

the joke he suggests that they have marijuana but it has been medically prescribed.

Similar to Extract 1, this extract demonstrates a conversational humor speech pattern that goes back and forth between participants and ends with a punchline. In this extract, the participant who provided the joke's motive was also the one to provide a punchline. Participants who were listening to this conversation provided minimal responses to demonstrate positive attention to the speakers, which demonstrates a sense of support and empathy, highlighting the collaborative nature of the humor.

Extract 3: Conversational Humor

The third extract presented in **Table 4** follows the same conversational sequence that has been observed in the previous two extracts: Motive-Joke-Joke return-Punchline. However, the difference in this extract is that rather than the collective joking that was observed in **Tables 2, 3**, the joke in this extract is directed at a specific participant (Victor). Where Extracts 1 and 2 presented collaborative humor and collective joking about the experiences of living with pain was demonstrated, in Extract 3 the joke is directed at another participant (Victor), initiating an intervention from a healthcare professional. This intervention functioned to ensure that no participants felt victimized by other participants and rather that the play frame of the conversation was mutually agreed upon.

From the extract, it is suggested that there is no malice intended. On the contrary, the teasing might be a form of

building relations and using jokes as a coping mechanism to deal with chronic pain. Victor provides the motive for the joke by discussing his swollen limb (Line 4). After Victor says, "Just pain killer is enough hhh" (Line 8) suggesting that his situation is not too serious, and a pain killer is enough to treat it, Tyrone initiates the joke: "Yeah he's got an extra leg to change over" (Line 9). The minimal response laughter by Helen and Tyrone in Lines 10 and 11 suggests that this conversation is within a play frame, and what has been said should be interpreted as a joke. Brenda returns the joke in line 12, and Tyrone continues this return in line 13. The punchline in this conversational joke is provided by Brenda in line 15: "Yeah yeah take it off=." Although the participants' jokes seem targeted toward Victor, it is noted that the joke initiation is made after Victor himself downplays the severity of his situation, followed with laughter and an exhale. The uptake of this motive and its conversion to a joke then demonstrates that the participants acknowledge that Victor is understating his situation and therefore they are using conversational humor to provide support and lighten the conversation. In this extract, the healthcare professional intervenes to ensure no malice is intended toward Victor (Lines 17–19 and 21, 22). Tyrone responds with "No no no no no" further confirming that the joke was not intended to harm Victor. An intervention was required here as the joke (directed at a specific individual) could have been interpreted by Victor as malicious. Clarifying that the conversation was within a play frame allowed Victor to conceptualize the joke as it was intended, rather than experiencing it as direct teasing. Therefore, although

TABLE 4 | Extract 3 Joking cycle.

Line number	Pseudonym	Conversation text	Conversation analysis
4	Victor:	Yeah uh my limb is swollen at the ↓moment	Set up of joke (Motive)
5	Brenda:	Really?	
6	Victor:	Yeah:	
7	Brenda:	Have you got anything for it?	
8	Victor:	Just pain killer is enough heh hhh	
9	Tyrone:	Yeah he's got an extra leg to change over	Joke 1 presented
10	Helen:	Hehhh	Minimal responses
11	Tyrone:	Hehh	
12	Brenda:	Send one over here then Victor when you find one, will you?	Joke return
13	Tyrone:	He's got two spare ones so when he get when he gets pain in	Joke 2 presented
14		one=	
15	Brenda:	=Yeah yeah take it off=	Joke punchline
16	Tyrone:	=Yeah	
17	HCP:	.h Look at this sense of humor who would have thought in	Intervention
18		Week 1 you would have been <u>laughing</u> about having (.)	
19		about having chronic pain. And here we are sat here:	
20	Brenda:	I know=	
21	HCP:	=I know it's great knowing it's meant as a <u>joke</u> there's no	
22		<u>malice</u> in it	
23	Tyrone:	No no no no no	Acknowledgment of play frame

HCP, Healthcare Professional.

the joke is directed at an individual, the collaborative joke was used to find positive aspects of their chronic pain experiences and lighten the collective mood.

Extracts 1–3 therefore demonstrate the stability of the first speech pattern: Motive-Joke-Joke return-Punchline, used for collaboratively for conversational humor. Using conversational humor was a sign of social support and a coping mechanism to help participants understand and make sense of their chronic pain experiences and maintain interactional levity, despite the seriousness of the pain conditions they all lived with.

The Venting Cycle

The second speech pattern observed in the data was a venting cycle in which patients listen to each other vent about their experiences of chronic pain and pain management. The sequence of conversational venting was begun by an individual initiating an extended outline of a personal frustration, venting initiation. This was then extended or escalated further and then a joke punchline was offered either by the person venting or by a colleague to complete the venting and facilitate a change of conversational topic. The joke punchline therefore served as a pivot in conversation whereby the person who held the floor would pass their turn over to someone else (37). After the turn was passed over, the expression of negative feelings passed from one participant to another or entirely stopped. The punchline also demonstrated conversational engagement among participants, which served as a form of social support to

help participants make sense of their experiences and validate their vent.

Extract 4: Venting Cycle

Extract 4 (shown in Table 5) presents a scenario where two patients converse about chemists and the difficulties of obtaining (correct) prescriptions. In this extract, both participants talk and listen to each other vent about their shared negative experiences of seeking pain medication:

Brenda initiates her venting in line 173 by describing her negative experience with a chemist. Steve interrupts and shares his common experience in Line 176, which demonstrates the continuation of the vent. Steve and Brenda continue to share their negative experiences between lines 179–184. In line 183, Brenda interrupts Steve with overlapping speech and resumes her vent, showing an upgraded agreement to his venting by sharing a related experience related to his situation (lines 185–187). From lines 188 to line 197, Steve provides minimal responses such as “Yeah” (Line 188), “Oh crikey no” (Line 190) and “No” (Line 194) in response to the escalating venting by Brenda (Lines 188, 190, 194, and 197). These minimal responses allow for the venting to continue as it provided positive reinforcement and attentional acknowledgment. Steve initiates the joke punchline in line 199: “Sometimes >take a kick up the bum and then< heh” and Brenda adds to the joke: “Just like she needs u:m fireworks.” Following the joke punchline, there is a conversational pivot from a mutual vent to a more general conversation. Additionally, a change in turn taking is observed from Brenda who was the

TABLE 5 | Extract 4: Venting cycle.

Line number	Pseudonym	Conversation text	Conversation analysis
173	Brenda:	Huh that's good because our chemist is absolutely (.)	Vent initiation
174		<u>Terrible</u> . (.) never get our medication on time: I had to ring	
175		up and complain about it=	
176	Steve:	=Yeah I was suffering from that °I was getting a lot of th	Venting continues
177		cos I was like the ones that was um: I actually ran out of	
178		<u>patches</u> and you know I can't do without them because-	
179	Brenda:	>Of course not<	
180	Steve:	Because you: <u>need</u> them for the one for the <u>pain</u> yes, but	
181		<u>also</u> there's that you get withdrawal symptoms if you're <u>not</u>	
182		taking them	
183	Brenda:	[In fact um-]	
184	Steve:	[They can't win though-]	
185	Brenda:	One time, I always used to say to the pharmacist the last (.)	Venting escalates
186		and when I'm down to two patches (.) >can you <u>order</u> them	
187		for me please<	
188	Steve:	Yeah	
189	Brenda:	I say because I don't wanna run out I'd rather-	
190	Steve:	Oh crikey <u>NO</u>	
191	Brenda:	I'd rather be 2 weeks I have 2 weeks supply >and then	Venting continues
192		all of a sudden< forget about it and run out but but um	
193		but um: she wasn't doing it? see	
194	Steve:	°No°	
195	Brenda:	So: my husband then complained to them about it and then	
196		the woman said oh I'll look into this and she <u>did</u>	
197	Steve:	Yeah	
198	Brenda:	And <u>now</u> the pharmacist is spot on	
199	Steve:	Sometimes >take a kick up the bum and then< heh	Joke punchline
200	Brenda:	Just like she needs u:m fireworks	
201	Steve:	Hehhhhh	Minimal response
202	Brenda	°So yeah°	
203	Steve:	Anyway	Change in turn

primary initiator of the venting, allowing Steve to take the floor. This extract demonstrates a Venting initiation-Continuation and escalation of venting-Joke punchline to finish the venting speech pattern. The mutual use of minimal responses allowed the individuals to continue to express their negative experiences to each other.

Extract 5: Venting Cycle

Demonstrating a similar speech pattern as Extract 4, the fifth extract presented in **Table 6** shows an interaction between two individuals who discuss pain medication and the worsening of their pain symptoms. In this extract, one participant releases their negative emotions while the other participant listens:

Steve initiates the vent by discussing the amount of medication he has to take: “When my wife comes back with her big bag of meds, and I go” (Line 22) “Is that all mine? and she goes yeah. oh bugger (.)” (Line 24). He discusses his “pain level

getting worse” and his pain escalation as a result of starting the medication at the wrong time, which demonstrates an escalation in his venting. Finally, Steve provides the joke punchline: “=So I’m going to be going through the alphabet anyways I’ll just Get through it ↑°hopefully°.” In this extract, Steve is the participant who is venting and he ends his own venting with a joke. Similar to the first extract, a conversational pivot occurs after the punchline as there is a change in turn taking and Brenda begins to share her experience regarding her prescribed medication.

In this extract, Brenda provides backchannel responses throughout the conversation. These include “Yeah” (Line 23), “=Uh huh uh huh=” (Line 27) and “Ah:=” (Line 34). This demonstrates that she is engaged and actively listening to Steve. Additionally, in line 42, Brenda asks, “What did you say what milligram you’re on,” which indicates engagement, positive politeness and curiosity as she has followed up with a question that was related to Steve’s expression of emotion. The Venting initiation-Continuation and escalation of venting-Joke punchline

TABLE 6 | Extract 5: Venting cycle.

Line number	Pseudonym	Conversation text	Conversation analysis
22	Steve:	When my wife comes back with her big bag of <u>meds</u> and I go.	Vent initiation
23	Brenda:	Yeah	
24	Steve:	Is that all <u>mine</u> ?and she goes yeah. oh bugger (.) I'm just going	
25		through a um change in my medication now so I'm	Venting continues
26		going through (.) quite a bit of problems with pain and stuff so=	
27	Brenda:	=Uh huh uh huh=	Minimal response
28	Steve:	=My pain level is getting worse ↑ I started	Venting escalates
29		>it a bit at the wrong time because<	
30		this <u>literally</u> this week is my first week on this new thing	
31	Brenda:	Uh huh=	
32	Steve:	=And I've got to <u>reduce</u> and <u>reduce</u> and reduce and then	
33		increase the other bits h	
34	Brenda:	Ah:=	
35	Steve:	=So I'm going to be going through the alphabet anyway I'll just	Joke punchline
36		get through it ↑"hopefully"	
37	Brenda:	Ok um: the doctor gave me medication prescribed 'em to me	Change in turn
38		this morning	
39	Steve:	Yeah	
40	Brenda:	He prescribed me [medication name] yeah	
41	Steve:	Well I've got them yeah:	
42	Brenda:	What did you say? what milligram you're on	
43	Steve:	I <u>used</u> to have two:	

to finish the venting is consistent in this extract with engagement and active listening by all participants.

Extract 6: Venting Cycle

The final extract, shown in **Table 7**, presents a situation where five patients have a conversation about pain medications and medical trials. Even though five members are present in the conversation, one participant vents about their experience with medical medicine trials whilst the others actively listen.

In suggesting that they do not “want to be treated as a guinea pig” (Line 133) for medical trials, Brenda initiates the venting. Immediately Steve provides a minimal response of ‘no’ (Line 134). This response indicates that there is engagement and active listening as he voices his opinion to agree with what Brenda proposed. Steve to express his negative experiences in lines 135, 136, and Gemma gives an upgraded agreement indicating collaborate support for to the context and content of the venting: “Yeah my surgery is the same it’s ridiculous” (Line 137). Brenda continues and escalates her venting about her medication and pain experiences from Line 138–169. Throughout this section, participants provide minimal responses and agreement to Brenda, who is holding the floor. These responses include laughter, “Yeah hehh” (Line 159) and minimal agreement such as “Well yeah” (Line 153), “Yeah” (Line 163), and “Mm” (Line 164). These responses suggest agreement, engagement and responsiveness from the participants listening to their colleague speak about their experience. Additionally, Tyrone repeats Brenda’s negative sentiment about patients being

guinea pigs in Line 175. This further demonstrates displays of agreement toward Brenda, which provides the incentive for Brenda to continue her vent.

Participants are demonstrating recognition of the need to move beyond the vent and therefore the joke punchline was initiated by Brenda, who began the vent: “Try this: try that:” (Line 176) and Tyrone adds to the punchline: “Try that like it it might take your pain away but like I say it might make you looney or something like that” (Line 177, 178). Again, the joke punchline pivots the conversation from a participant venting to Tyrone taking the floor and speaking about his experience in comparison to Brenda’s experience. This is acknowledged with a minimal “hehh” (Line 181) by Gemma, which demonstrates the ongoing play frame of the conversation.

The venting cycle extracts have demonstrated that pain venting cycles reliably occur among patient-to-patient interactions. The speech pattern observed in venting cycles is as follows: Vent initiation-Continuation of venting-Escalation of venting-Joke punchline, but it can be both collaborative venting, or venting by a single person, supported by others. Actively listening to other participants’ venting adds to the sense of shared pain experience and allows for an individual’s feelings to be validated by individuals with similar experiences.

DISCUSSION

Results from this observational study demonstrated that patient-to-patient conversations within pain management settings were

1027 **TABLE 7** | Extract 6: Venting cycle.

Line number	Pseudonym	Conversation text	Conversation analysis
1031	Brenda:	I've told the doctor I don't >want to be treated as guinea pig<	Vent initiation
1032	Steve:	No h	
1033		right try me on this one: try me on <u>that</u> one: no (,) I said no	Venting continues
1034		want the proper medication	
1035	Gemma:	°Yeah my surgery is the same it's ridiculous°	
1036	Brenda:	And >once the other day going to the doctor< and he said t	Venting escalates
1037		me oh. you've got to come of this tablet you can't have it	
1038		anymore and uh and my husband said (,) hang on you've	
1039		just got the medication right! you can't just say to her	
1040		mo she's not gonna have it=it's like he gave me morphine	
1041		I've passed out with morphine patches it started off (,) With	
1042		the 12 and then I've went up to 75.h and um he said to me	
1043		I might have to take you off them and I said NO YOU WON'T	
1044		I said I'm not going to be <u>crawling in agony</u>	
1045			
1046	Steve:	No	
1047	Brenda:	And he said to me=my doctor said to me I'm gonna try	Venting continues
1048		on um morphine tablets ↓oh my God. the first week I was	
1049		supposed to take a tablet in the evening the second week	
1050		I'm supposed to take <u>two</u> tablets on the first week I was <u>sick</u>	
1051		<u>all</u> : week with them	
1052	Steve:	Well yeah	
1053	Brenda:	And then my ankle from here down <u>blew</u> up like that	
1054		I couldn't get off the <u>settee</u> I couldn't my husband	
1055		couldn't even help me to get off and I saw the doctor about it	
1056		he said I want you to <u>stop</u> them <u>immediately</u> and I said	
1057		I did. Don't you worry	
1058			
1059	Steve:	[Yeah hehh]	
1060	Gemma:	[Hehh]	
1061		To touch them you are <u>lucky</u> to be al to be <u>alive</u>	
1062	Brenda:	I said ! <u>TOOK</u> what <u>YOU</u> <u>GAVE</u> me	
1063	Gemma:	Yeah	
1064	Steve:	Mm	
1065	Brenda:	And I said to him I am <u>not</u> a <u>guinea</u> pig and apparently the	Venting escalates
1066		drugs he gave me were for (,) depression and epileptic fits	
1067		which I don't suffer with (,) there are two: two medications	
1068		into one and I don't even suffer from epileptic fits <u>or</u>	
1069		depression	
1070			
1071	Gemma:	Yeah h	
1072	Steve:	.h No	
1073	Kate:	So: I take it that it-	
1074	Tyrone:	With the doctor I'd-	
1075	Brenda:	See! so what's the point!	
1076	Tyrone:	Patients are guinea pigs >you know what I ↑mean<	Venting continues
1077	Brenda:	[Try thi:s try tha:t]	Joke punchline
1078	Tyrone:	[Try that like] it it might take your pain away but like I say	Joke punchline
1079		it might make you <u>looney</u> or something like that	Joke punchline
1080	Steve:	°It's it's a side effect°	Change in turn
1081	Tyrone:	.h It's like that when they gave <u>me</u> amitriptyline	

1141 employing two specific interactional mechanisms, identified
 1142 via conversation analysis. These approaches to communication
 1143 were: (1) conversational humor and (2) a venting cycle. These
 1144 patterns in speech demonstrated how CP patients expressed
 1145 their pain experiences and supported each other through
 1146 constructive interpersonal interactions. Both conversational
 1147 mechanisms involved the use of humor, collaborating together
 1148 to describe pain experiences with levity. These conversations
 1149 allowed PLwCP space to express their negative emotions, before
 1150 ending this cathartic vent with a light-hearted punchline, which
 1151 served to pivot the conversation toward other contributors
 1152 and new topics. Although the humor was targeting difficult
 1153 experiences of pain and pain management, the positive reception
 1154 and adoption of humor within the group indicated that the
 1155 humor facilitated the discussion of sensitive topics in a socially
 1156 supportive environment.

1157 The results of this study showed how fundamental the use
 1158 of humor is in interpersonal interactions, even within a pain
 1159 management setting. The deliberate application of humor to
 1160 difficult situations has been demonstrated in other chronic
 1161 illness populations; after radiotherapy, humor enabled people
 1162 with cancer to cope better with their health (38). Similarly,
 1163 quantitative assessment of humor use within pain, has suggested
 1164 that PLwCP may employ humor to gain perspective on their
 1165 pain and distance themselves from their situation by accepting
 1166 the negative emotions that common when living with chronic
 1167 pain (39). In confirmation, a recent narrative review on pain and
 1168 humor suggested that humor serves to improve pain coping and
 1169 minimize emotional distress (40). Humor in pain management
 1170 settings may therefore act as a method of bolstering pain-related
 1171 resilience and reducing pain catastrophising (41). Certainly, in
 1172 this study, the deliberate use of humor was evidenced through
 1173 consistent use of jokes and delivery of punchlines, even when
 1174 these may be least expected, for example, after venting and
 1175 describing difficult pain-related situations. Finding the humor
 1176 in chronic pain is indicative of highly subjective, black humor,
 1177 which is mutually understood and facilitated by participants'
 1178 common experiences, but which may not fully translate to
 1179 others without pain or an understanding of chronic pain. So-
 1180 called "black humor" or "gallows humor" is recognized as
 1181 having therapeutic value, particularly when used by individuals
 1182 facing trauma or challenging circumstances in health settings
 1183 (42). Indeed, in clinical contexts, such self-deprecating humor
 1184 has been found to boost psychological well-being, despite its
 1185 dark(er) content (43), suggesting it is adaptive in the pain
 1186 management context.

1187 The theoretical underpinning of the humor evidenced
 1188 in the current study provides an interesting disconnect
 1189 between our findings and previous interpretations of humor
 1190 in pain management. Previous research, using a cross-sectional
 1191 quantitative assessment of humor, pain and psychological well-
 1192 being has been interpreted as indicating that humor is a
 1193 distancing mechanism, a way of making space between chronic
 1194 pain and oneself as a method of self-protection (39). However,
 1195 the real-life conversational mechanisms demonstrated in this
 1196 study did not suggest that interactive humor was a distancing
 1197 mechanism. Indeed, an approach-based pattern was found; in

1198 the naturalistic interactions we recorded, there was a clear
 1199 willingness to "vent" by giving detailed descriptions about
 1200 one's pain experience, setting up a conversational context in
 1201 which PMP group members actively chose to *enter into* the
 1202 discussion and to collaborate in joking about pain. This suggests
 1203 a willingness to approach rather than avoid the pain situation that
 1204 is described by their colleague and a willingness to approach one's
 1205 own pain when venting.

1206 The difference between our findings here and that of Ramírez-
 1207 Maestre et al. (39) may be explained by the "optimal matching
 1208 hypothesis" (9). In the current study, all participants already had
 1209 a mutual understanding of chronic pain and therefore the shared
 1210 pain experience was a commonality between the group, allowing
 1211 for psychological safety in approaching this difficult topic (44)
 1212 removing the need to employ distancing. Therefore, patient-to-
 1213 patient interactions in PMPs were evidencing an *affiliative humor*
 1214 *style*, using humor to enhance one's relationship with others
 1215 (45). This is powerful evidence that interpersonal, conversational
 1216 humor in PLwCP is not aggressive, self-enhancing, self-defeating
 1217 or distancing. It is constructive, collaborative, communicative
 1218 and creates warm appreciation of others within the group,
 1219 respecting the shared mutual understanding of chronic pain.
 1220 Further research is needed to establish whether this humor style
 1221 only worked because of group dynamics or whether it can be
 1222 identified more widely. Where "dark humor" is employed, there
 1223 is the risk that such humor could be misconstrued, causing
 1224 unease, alienation or even perceived insult (46). The participants
 1225 in this study were all PLwCP, which provided a certain level of
 1226 implicit understanding of pain experiences that is likely not there
 1227 in interactions between different patient populations or between
 1228 people with and without CP.

1229 The prosocial nature of the humor demonstrated in this
 1230 study was unusual in that it employed mutual sarcasm, with
 1231 the sarcasm applied not to a person, but to the context under
 1232 discussion—chronic pain itself. Wider literature suggests that the
 1233 type of humor is important when addressing health conditions
 1234 (47, 48) for example, mocking and sarcasm are typically viewed
 1235 negatively (38). However, in this study, the participants used
 1236 sarcasm that was received positively and was directed toward
 1237 pain rather than a person or their behavior. The sarcasm
 1238 highlighted the ridiculousness of the situation (for example,
 1239 being prescribed numerous medications) rather than ridiculing
 1240 the speaker, suggesting that the target of the humor may be
 1241 more important compared to humor type in interactions between
 1242 PLwCP. The current study demonstrated one incidence (Extract
 1243 3) when use of sarcasm came closer toward being associated with
 1244 an individual, rather than pain more generally. In that instance,
 1245 the healthcare professional stepped in to maintain the prosocial
 1246 balance in the humor, though the participants in the conversation
 1247 showed strong awareness of the need to keep their conversational
 1248 humor inclusive. It is possible therefore that HCPs may benefit
 1249 from additional awareness and training in the value of humor
 1250 within the pain management setting, to respond to and encourage
 1251 such humor effectively.

1252 The venting cycle that was demonstrated in this research
 1253 found that PLwCP were active listeners to their colleagues'
 1254 distress, using positive minimal response (mmm, yeah) and

interjecting shared experiences to demonstrate empathetic listening. This shows that group interactions in the PMP setting are building pain coping, as has also been shown in chronic pain support groups (41). Yet, the venting and the humor which concludes the vent cycle are not a distraction away from pain, posited as a primary theoretical explanation for the benefits of humor in pain management (40), nor are they methods of promoting cognitive reappraisal of pain (49). Instead, the venting cycle, is further evidence of the power of social support to improve pain coping and pain management (50). The vent is allowed to continue by the listeners, who help the speaker through the expression of their difficult experience. Therefore, the vent is acting as a method of emotion regulation, which can help to mediate the relationship between negative emotions and anxiety or depressive disorders (51).

Limitations and Future Recommendations

The recruitment for the current study was only from one of the three potential PMPs, and participants were recruited from one geographical site. Whilst this may limit generalizability of the findings, this is the first study, to the authors' knowledge, that uses CA to examine conversations between CP patients rather than between a health professional and CP patients. Significant efforts were made to ensure that naturalistic conversation could be recorded, but the presence of audio-visual recording equipment inevitably limits this. Future research could aim to hold a focus group with participants after the conversational mechanisms are identified via CA, to allow participants to validate the observations. Future research could aim to confirm with participants *post-hoc* the reasoning for the conversational humor and venting cycle. The findings need to be replicated in comparable settings and confirmed *post-hoc* with CP patients before the universality of such mechanisms can be reliably confirmed. Additionally, the cultural context of the current study was within a British, Caucasian sample, and it is likely that individuals from other cultures may not employ or perceive humor as a positive conversation mechanism which nurtures social support. It is possible that without a common understanding of chronic pain, pain humor could be misperceived as exclusionary or an insult (46), which would jeopardize group cohesion and clinical outcomes. Future research could assess patient-to-patient interactions with a more culturally diverse sample, and across a variety of inpatient and outpatient pain management settings to identify commonalities and variation in pain humor usage.

REFERENCES

- Briggs AM, Woolf AD, Dreinhöfer K, Homb N, Hoy DG, Kopansky-Giles D, et al. Reducing the global burden of musculoskeletal conditions. *Bull World Health Organ.* (2018) 96:366–8. doi: 10.2471/BLT.17.204891
- Ellis B, Ly M, Steinberger S. *Unseen, Unequal and Unfair: Chronic Pain in England.* Versus Arthritis (2021). Available online at: <https://www.versusarthritis.org/news/2021/june/unseen-unequal-and-unfair-chronic-pain-in-england/> (accessed March 28, 2022).
- Froud R, Amundsen PA, Bartys S, Battie M, Burton K, Foster NE, et al. Opportunities and challenges around adapting supported

CONCLUSIONS

Humorous interpersonal interactions about chronic pain act as an important forum for social support-building between members of a PMP. Such dynamic interactions referenced the mutual experience of chronic pain and fostered common humor, helping people living with chronic pain to make sense of their pain in a prosocial atmosphere. When negative emotions were strongly expressed through venting, responses were empathetic with active listening used to validate the negative emotions, allowing for a cathartic expression of personal challenges. Patient-to-patient interactions within the PMP were therefore strongly prosocial and mutually inclusive, evidencing that the socially communicative context fostered during PMPs has significant power to facilitate enhanced clinical outcomes and group cohesion.

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by NHS Ethics REC: 13/WM/0214. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

KF and SP: conceptualization of research. KF: data collection. KF, KA, and AM: data analysis and drafting the manuscript. KF, KA, AM, and SP: editing and review of manuscript. All authors contributed to the article and approved the submitted version.

ACKNOWLEDGMENTS

The authors would like to acknowledge the support of Alena Meister who assisted in data collection and Milton Keynes University Hospital Department of Pain Management who provided the setting for the research.

- employment interventions for people with chronic low back pain: modified nominal group technique. *Disabil Rehabil.* (2021) 43:2750–7. doi: 10.1080/09638288.2020.1716863
- Solé E, Racine M, Tomé-Pires C, Galán S, Jensen MP, Miró J. Social factors, disability, and depressive symptoms in adults with chronic pain. *Clin J Pain.* (2020) 36:371–8. doi: 10.1097/AJP.0000000000000815
 - Dueñas M, Ojeda B, Salazar A, Mico JA, Failde I. A review of chronic pain impact on patients, their social environment and the health care system. *J Pain Res.* (2016) 9:457–67. doi: 10.2147/JPR.S105892
 - Jensen MP, Ehde DM, Hoffman AJ, Patterson DR, Czerniecki JM, Robinson LR. Cognitions, coping and social environment

- 1369 predict adjustment to phantom limb pain. *Pain*. (2002) 95:133–
1370 42. doi: 10.1016/S0304-3959(01)00390-6
- 1371 7. Chang PF, Bazarova NN, Wethington E. How older adults with
1372 chronic pain manage social support interactions with mobile media.
1373 *Health Commun*. (2020) 37:1–13. doi: 10.1080/10410236.2020.184
1374 6272
- 1375 8. Rising CJ, Bol N, Burke-Garcia A, Rains S, Wright KB. Perceived stress
1376 in online prostate cancer community participants: examining relationships
1377 with stigmatization, social support network preference, and social support
1378 seeking. *J Health Commun*. (2017) 22:469–76. doi: 10.1080/10810730.2017.13
1379 04471
- 1380 9. Cutrona CE, Russell DW. Type of social support and specific stress: toward a
1381 theory of optimal matching. In: Sarason BR, Sarason IG, Pierce GR, editors.
1382 *Social Support: An Interactional View*. New York: John Wiley and Sons (1990).
1383 pp. 319–66.
- 1384 10. Seawell AH, Cutrona CE, Russell DW. The effects of general social support
1385 and social support for racial discrimination on African American women's
1386 well-being. *J Black Psychol*. (2014) 40:3–26. doi: 10.1177/009579841246
1387 9227
- 1388 11. Groth M, Grandey A. From bad to worse: negative exchange spirals in
1389 employee–customer service interactions. *Organ Psychol Rev*. (2012) 2:208–
1390 33. doi: 10.1177/2041386612441735
- 1391 12. Barsade SG. The ripple effect: emotional contagion and its influence on group
1392 behavior. *Administr Sci Q*. (2002) 47:644–75. doi: 10.2307/3094912
- 1393 13. Hatfield E, Rapson R. Love and intimacy. *Encyclopedia Hum Behav*.
1394 (1994) 3:93–101.
- 1395 14. Simm R, Barker C. Five years of a community pain service solution-focused
1396 pain management programme: extended data and reflections. *Br J Pain*. (2018)
1397 12:113–21. doi: 10.1177/2049463717744358
- 1398 15. Hepburn A, Hepburn DA, Wiggins S. *Discursive Research in Practice:
1399 New Approaches to Psychology and Interaction*. New York, NY: Cambridge
1400 University Press (2007).
- 1401 16. Wittgenstein L. *Philosophical Investigations* (Translated by GE Anscombe).
1402 Chichester: Blackwell (1953).
- 1403 17. Heritage J. Conversation analysis: methodological aspects. In: *Aspects of
1404 Oral Communication*. De Gruyter (2011). pp. 391–418. Available online
1405 at: [https://www.degruyter.com/document/doi/10.1515/9783110879032.391/
1406 html](https://www.degruyter.com/document/doi/10.1515/9783110879032.391/html) (accessed March 28, 2022).
- 1407 18. Peräkylä A. Conversation analysis and psychoanalysis: Interpretation, affect,
1408 and intersubjectivity. In: Peräkylä A, Antaki C, Vehviläinen S, Leudar I,
1409 editors. *Conversation Analysis and Psychotherapy*. Cambridge: Cambridge
1410 University Press (2008). p. 100–19. doi: 10.1017/CBO9780511490002.
1411 007
- 1412 19. Beach WA. Making cancer visible: unmasking patients' subjective experiences.
1413 *Health Commun*. (2019) 34:1683–96. doi: 10.1080/10410236.2018.15
1414 36941
- 1415 20. Peräkylä A. Conversation analysis: a new model of research in doctor–
1416 patient communication. *J R Soc Med*. (1997) 90:205–
1417 8. doi: 10.1177/014107689709000406
- 1418 21. Robinson JD, Heritage J. Intervening with conversation analysis: the case
1419 of medicine. *Res Lang Soc Interact*. (2014) 47:201–
1420 18. doi: 10.1080/08351813.2014.925658
- 1421 22. Stivers T, Timmermans S. Medical authority under siege: how clinicians
1422 transform patient resistance into acceptance. *J Health Soc Behav*. (2020)
1423 61:60–78. doi: 10.1177/0022146520902740
- 1424 23. Pilnick A, Trusson D, Beeke S, O'Brien R, Goldberg S, Harwood
1425 RH. Using conversation analysis to inform role play and simulated
1426 interaction in communications skills training for healthcare professionals:
1427 identifying avenues for further development through a scoping
1428 review. *BMC Med Educ*. (2018) 18:267. doi: 10.1186/s12909-018-
1429 1381-1
- 1430 24. Speer SA, McPhillips R. Initiating discussions about weight in a non-
1431 weight-specific setting: what can we learn about the interactional
1432 consequences of different communication practices from an
1433 examination of clinical consultations? *Br J Health Psychol*. (2018)
1434 23:888–907. doi: 10.1111/bjhp.12322
- 1435 25. Mace S, Collins S, Speer S. Talking about breast symmetry in the
1436 breast cancer clinic: what can we learn from an examination of
1437 clinical interaction? *Health Expect*. (2021) 24:209–21. doi: 10.1111/hex.
1438 13144
- 1439 26. British Pain Society. Guidelines for Pain Management Programmes for
1440 Adults. London: The British Pain Society. (2013). Available online at: [https://
1441 www.britishtainsociety.org/static/uploads/resources/files/pmp2013_main_
1442 FINAL_v6.pdf](https://www.britishtainsociety.org/static/uploads/resources/files/pmp2013_main_FINAL_v6.pdf) (accessed March 28, 2022).
- 1443 27. Jefferson G. Glossary of transcript symbols with an introduction. In: Lerner G,
1444 editor. *Conversation Analysis. Studies From the First Generation*. Amsterdam;
1445 Philadelphia, PA: Benjamins (2004). pp. 13–31.
- 1446 28. Ide N, Pustejovsky J, editors. *Handbook of Linguistic Annotation*. Dordrecht:
1447 Springer Netherlands (2017).
- 1448 29. Plumridge E, Goodyear-Smith F, Ross J. Nurse and parent
1449 partnership during children's vaccinations: a conversation analysis.
1450 *J Adv Nurs*. (2009) 65:1187–94. doi: 10.1111/j.1365-2648.2009.
1451 04999.x
- 1452 30. Schegloff EA. *Sequence Organization in Interaction: A Primer in Conversation
1453 Analysis I*. Cambridge: Cambridge University Press (2007).
- 1454 31. Peräkylä, A. (2011). Validity in research on naturally occurring social
1455 interaction. In D. Silverman (Ed.), *Qualitative Research*. Third Edition. (3rd
1456 ed.). Sage: London.
- 1457 32. Bateson G. *A Theory of Play and Fantasy*. Cambridge, MA: MIT Press (1972).
- 1458 33. Brown P, Levinson SC. Universals in language usage: politeness phenomena.
1459 In: Goody EN, editor. *Questions and Politeness: Strategies in Social Interaction*.
1460 Cambridge: Cambridge University Press (1978). pp. 56–311.
- 1461 34. Kitamura N. Adapting brown and levinson's 'politeness' theory to the analysis
1462 of casual conversation. In: *Proceedings of ALS2k, the 2000 Conference of
1463 the Australian Linguistic Society*. (2000). p. 1–8. Available online at: [https://
1464 d1wqtxts1xzle7.cloudfront.net/52385870/kitamura-with-cover-page-v2.pdf?
1465 Expires=1648474523&Signature=Osv-7wcj0s0EW~GH-jlr1yhLY4KdDFK~
1466 XcU9GvtXoLMjkVNNwdR~fhLJDVYsjrtWRcqYhQ6oS3t01QXjG8s~
1467 BfQTE748SAmF3a78-BV~Ols9-X3XWVjHekia9tMG0PmlC~
1468 D69hMJMrRTn779OgzyqV4fjPcnV-XfBVxIYpkZf~QLsArHK~
1469 JvqCDvfmQul~aWg7823ZrwlRLUpbW8zwYFjX-mWxfLZu8hUPKkop~
1470 rgab6x71BuzSmgXq2zhZdpfjgTlQwzOkIPWUuoc6GEUHCmcdcsVHjkQNW~
1471 Aswg6LTT37Wm6UZU~O9Ic795eSKm5OibFL2lBGYP04YFnse0cQ__&
1472 Key-Pair-Id=APKAJLOHF5GGSLRBV4ZA](https://d1wqtxts1xzle7.cloudfront.net/52385870/kitamura-with-cover-page-v2.pdf?Expires=1648474523&Signature=Osv-7wcj0s0EW~GH-jlr1yhLY4KdDFK~XcU9GvtXoLMjkVNNwdR~fhLJDVYsjrtWRcqYhQ6oS3t01QXjG8s~BfQTE748SAmF3a78-BV~Ols9-X3XWVjHekia9tMG0PmlC~D69hMJMrRTn779OgzyqV4fjPcnV-XfBVxIYpkZf~QLsArHK~JvqCDvfmQul~aWg7823ZrwlRLUpbW8zwYFjX-mWxfLZu8hUPKkop~rgab6x71BuzSmgXq2zhZdpfjgTlQwzOkIPWUuoc6GEUHCmcdcsVHjkQNW~Aswg6LTT37Wm6UZU~O9Ic795eSKm5OibFL2lBGYP04YFnse0cQ__&Key-Pair-Id=APKAJLOHF5GGSLRBV4ZA) (accessed March 28, 2022).
- 1473 35. Pomerantz A. *Second Assessments: A Study of Some Features of Agreements*.
1474 Disagreements. Irvine, CA: University of California (1975).
- 1475 36. Coates J. *Women, Men and Language: A Sociolinguistic Account of
1476 Gender Differences in Language*. 3rd ed. New York, NY: Routledge (2015).
1477 doi: 10.4324/9781315645612
- 1478 37. Goffman E. *Frame Analysis: An Essay on the Organization of Experience*.
1479 Cambridge, MA: Harvard University Press (1974). pp. ix, 586.
- 1480 38. Samant R, Balchin K, Cisa-Paré E, Renaud J, Bunch L, McNeil A,
1481 et al. The importance of humour in oncology: a survey of patients
1482 undergoing radiotherapy. *Curr Oncol*. (2020) 27:350–3. doi: 10.3747/co.27.
1483 5875
- 1484 39. Ramírez-Maestre C, Esteve R, López-Martínez AE, Miró J, Jensen MP, de la
1485 Vega R. Beyond pain intensity and catastrophizing: the association between
1486 self-enhancing humour style and the adaptation of individuals with chronic
1487 pain. *Eur J Pain*. (2020) 24:1357–67. doi: 10.1002/ejp.1583
- 1488 40. Pérez-Aranda A, Hofmann J, Feliu-Soler A, Ramírez-Maestre C, Andrés-
1489 Rodríguez L, Ruch W, et al. Laughing away the pain: a narrative review
1490 of humour, sense of humour and pain. *Eur J Pain*. (2019) 23:220–
1491 33. doi: 10.1002/ejp.1309
- 1492 41. Finlay KA, Peacock S, Elander J. Developing successful
1493 social support: an interpretative phenomenological analysis of
1494 mechanisms and processes in a chronic pain support group.
1495 *Psychol Health*. (2018) 33:846–71. doi: 10.1080/08870446.2017.142
1496 1188
- 1497 42. Pérez-Aranda A, García-Campayo J, Gude F, Luciano JV, Feliu-Soler A,
1498 González-Quintela A, et al. Impact of mindfulness and self-compassion on
1499 anxiety and depression: The mediating role of resilience. *Int J Clin Health
1500 Psychol*. (2021) 21:100229. doi: 10.1016/j.ijchp.2021.100229
- 1501 43. Navarro-Carrillo G, Torres-Marín J, Corbacho-Lobato JM, Carretero-Dios H.
1502 The effect of humour on nursing professionals' psychological well-being goes
1503 beyond the influence of empathy: a cross-sectional study. *Scand J Caring Sci*.
1504 (2020) 34:474–83. doi: 10.1111/scs.12751

1483 44. Grailey KE, Murray E, Reader T, Brett SJ. The presence and potential
 1484 impact of psychological safety in the healthcare setting: an evidence
 1485 synthesis. *BMC Health Serv Res.* (2021) 21:773. doi: 10.1186/s12913-021-
 1486 06740-6
 1487 45. Martin RA, Puhlik-Doris P, Larsen G, Gray J, Weir K. Individual
 1488 differences in uses of humor and their relation to psychological
 1489 well-being: development of the Humor Styles Questionnaire.
 1490 *J Res Pers.* (2003) 37:48–75. doi: 10.1016/S0092-6566(02)0
 1491 0534-2
 1492 46. Murphy SP. Humor orgies as ritual insult: putdowns and solidarity
 1493 maintenance in a corner donut shop. *J Contemp Ethnogr.* (2017) 46:108–
 1494 32. doi:10.1177/0891241615605218
 1495 47. Buiting HM, de Bree R, Brom L, Mack JW, van den Brekel
 1496 MWM. Humour and laughing in patients with prolonged incurable
 1497 cancer: an ethnographic study in a comprehensive cancer centre.
 1498 *Qual Life Res.* (2020) 29:2425–34. doi: 10.1007/s11136-020-
 1499 02490-w
 1500 48. Penson RT, Partridge RA, Rudd P, Seiden MV, Nelson JE,
 1501 Chabner BA, et al. Update: laughter: the best medicine?
 1502 *Oncologist.* (2005) 10:651–60. doi: 10.1634/theoncologist.10-
 1503 8-651
 1504 49. Papousek I, Rominger C, Weiss EM, Perchtold CM, Fink A, Feyaerts K.
 1505 Humor creation during efforts to find humorous cognitive reappraisals
 1506 of threatening situations. *Curr Psychol.* (2019) doi: 10.1007/s12144-019-
 1507 00296-9
 1508 50. Che X, Cash R, Ng SK, Fitzgerald P, Fitzgibbon BM. A
 1509 systematic review of the processes underlying the main and the
 1510 buffering effect of social support on the experience of pain. 1540
 1511 *Clin J Pain.* (2018) 34:1061–76. doi: 10.1097/AJP.0000000000000624 1541
 1512 51. Marr NS, Zainal NH, Newman MG. Focus on and venting of
 1513 negative emotion mediates the 18-year bi-directional relations
 1514 between major depressive disorder and generalized anxiety disorder
 1515 diagnoses. *J Affect Disord.* (2022) 303:10–17. doi: 10.1016/j.jad.2022.
 1516 01.079 1546
 1517 **Conflict of Interest:** The authors declare that the research was conducted in the
 1518 absence of any commercial or financial relationships that could be construed as a
 1519 potential conflict of interest. 1550
 1520 **Publisher’s Note:** All claims expressed in this article are solely those of the authors
 1521 and do not necessarily represent those of their affiliated organizations, or those of
 1522 the publisher, the editors and the reviewers. Any product that may be evaluated in
 1523 this article, or claim that may be made by its manufacturer, is not guaranteed or
 1524 endorsed by the publisher. 1555
 1525 *Copyright © 2022 Finlay, Madhani, Anil and Peacock. This is an open-access article
 1526 distributed under the terms of the Creative Commons Attribution License (CC BY).
 1527 The use, distribution or reproduction in other forums is permitted, provided the
 1528 original author(s) and the copyright owner(s) are credited and that the original
 1529 publication in this journal is cited, in accordance with accepted academic practice.
 1530 No use, distribution or reproduction is permitted which does not comply with these
 1531 terms.* 1560
 1532 1561
 1533 1562
 1534 1563
 1535 1564
 1536 1565
 1537 1566
 1538 1567
 1539 1568
 1540 1569
 1541 1570
 1542 1571
 1543 1572
 1544 1573
 1545 1574
 1546 1575
 1547 1576
 1548 1577
 1549 1578
 1550 1579
 1551 1580
 1552 1581
 1553 1582
 1554 1583
 1555 1584
 1556 1585
 1557 1586
 1558 1587
 1559 1588
 1560 1589
 1561 1590
 1562 1591
 1563 1592
 1564 1593
 1565 1594
 1566 1595
 1567 1596