



PEARL

**Estimating the number of people living with dementia at different stages of the condition in India: A Delphi process**

Farina, Nicolas; Rajagopalan, Jayeeta; Alladi, Suvarna; Ibnidris, Aliaa; Ferri, Cleusa P.; Knapp, Martin; Comas-Herrera, Adelina

**Published in:**  
Dementia

**DOI:**  
[10.1177/14713012231181627](https://doi.org/10.1177/14713012231181627)

**Publication date:**  
2023

**Link:**  
[Link to publication in PEARL](#)

**Citation for published version (APA):**

Farina, N., Rajagopalan, J., Alladi, S., Ibnidris, A., Ferri, C. P., Knapp, M., & Comas-Herrera, A. (2023). Estimating the number of people living with dementia at different stages of the condition in India: A Delphi process. *Dementia*, 0(0).  
<https://doi.org/10.1177/14713012231181627>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

**Estimating the number of people living with dementia at different stages of the  
condition in India: A Delphi process**

Nicolas Farina\*, <https://orcid.org/0000-0002-0635-2547>

Faculty of Health, University of Plymouth, Plymouth, UK

Jayeeta Rajagopalan, <https://orcid.org/0000-0002-1442-9786>

Strengthening Responses to Dementia in Developing Countries (STRiDE) India, National Institute of Mental Health and Neuro-Sciences (NIMHANS), Bangalore, India.

Suvarna Alladi, <https://orcid.org/0000-0002-0372-9572>

Strengthening Responses to Dementia in Developing Countries (STRiDE) India, National Institute of Mental Health and Neuro-Sciences (NIMHANS), Bangalore, India.

Aliaa Ibnidris, Department of Psychiatry and Mental Health, University of Cape Town, South Africa

Cleusa P Ferri, <https://orcid.org/0000-0002-1815-7685>

Department of Psychiatry, School of Medicine, Universidade Federal de São Paulo, Sao Paulo, Brazil

Martin Knapp, London School of Economics and Political Sciences, London, UK

Adelina Comas-Herrera, <https://orcid.org/0000-0002-9860-9062>

London School of Economics and Political Sciences, London, UK

\*Corresponding author.

This manuscript was accepted for publication on 19-May-2023 in *Dementia: the international journal of social research and practice*.

## **Abstract**

**Introduction:** Numerous studies have previously estimated the dementia prevalence in India. However, as these estimates use different methodologies and sampling strategies, generating definitive prevalence estimates can be difficult.

**Methods:** A Delphi process involving eight clinical and academic experts provided prevalence estimates of dementia within India, split by sex and age. The experts were also asked to estimate the number of people potentially living at different stages of the condition. *A priori* criteria were used to ascertain the point in which consensus was achieved.

**Results:** Our consensus estimates generated a dementia prevalence of 2.8% (95% CI = 1.9 to 3.6) for those aged 60 years and above in India. Consensus was achieved across age and sex prevalence estimates, with the exception of one (females aged 60-64). Our experts estimated that 42.9% of people living with dementia in India had a mild severity.

**Conclusions:** The findings indicate that there could be approximately 3.9 million people living with dementia in India, of which 1.7 million could be living with dementia of mild severity. Such estimates can better help researchers and policy makers to estimate the true cost and impact of dementia in India and can inform resource allocation decisions.

**Keywords:** Alzheimer's disease, diagnosis, severity, consensus, delphi

## **Introduction**

Population projections estimate that there are currently nearly 138 million older adults (aged 60+) in India, this is set to rise to almost 194 million by 2031 (National Commission on Population, 2020). Dementia is a condition that disproportionately affects older adults, leading to cognitive and functional decline. Estimating the numbers of people with dementia within a given country is essential to better understand the impact of dementia on a societal level. Within India, prevalence studies of dementia have been conducted across different regions of the country (e.g. Banerjee et al., 2017; Chandra et al., 1998; Das et al., 2006, 2008; Mathuranath et al., 2010; Poddar et al., 2011; Raina et al., 2008, 2014, 2016; Rajkumar et al., 1997; Razdan et al., 1994; Rodriguez et al., 2008; Saldanha et al., 2010; Seby et al., 2011; Shaji et al., 2005; Vas et al., 2001) with varying estimates generated. These differences in estimates may be attributed to variability in sociodemographic, cultural, genetic and environmental factors in addition to varying methodological approaches adopted by individual studies (Das et al., 2012). Through the use of statistical modelling, the Global Burden of Disease [GBD] India study estimated 3.69 million (95% UI 3.13–4.25) people were living with Alzheimer’s disease and other dementias in India in 2019 (Singh et al., 2021).

Meta-analysis can assist in synthesising and compiling these data from individual studies. Pooled prevalence estimates for dementia in India have ranged from 3.4% (95% CI = 2.0 to 5.0) to 4.4% (2.2-7.2) (Dhiman et al., 2021; Farina et al, 2020). However, a major factor to consider is that estimates are dependent on review inclusion criteria, which may be influenced by a whole range of methodological heterogeneity, most notably diagnostic criteria. For example, prevalence estimates have been found to be lower in those utilising DSM-IV diagnostic criteria in comparison to those using other diagnostic algorithms such as 10/66 diagnostic schedule (Farina et al., 2020; Paddick et al., 2013). This may be because the DSM-IV criteria are missing milder cases of dementia, or that the 10/66 algorithm is over-identifying cases of dementia (Paddick et al., 2013; Rodriguez et al., 2008). As a result of differences between prevalence estimates and individual study reporting styles, it can be difficult for systematic reviews to extract data to obtain more granular estimates (e.g., age, sex) in a meta-analysis. For example, very few systematic reviews (e.g., Choudhary et al., 2021) from India have provided prevalence of dementia split by age and sex, despite individual studies reporting on them. In addition, there are also considerable gaps in the existing evidence base, which makes it difficult to compile such data. A Delphi process is one approach that can help

overcome such gaps, through the use of experts to critically generate estimates using best evidence, whilst being able to draw from their expertise. Such approaches have been previously used to estimate dementia prevalence globally (e.g., Ferri et al., 2005).

Dementia is largely underdiagnosed, with only 1 in 10 persons with dementia receiving a diagnosis, treatment or care in India (Nulkar et al., 2019). Lack of awareness, stigma associated with the condition, shortage of resources such as health infrastructure and specialists to diagnose dementia contribute to this gap (Alzheimer's and Related Disorders Society of India, 2010). Consequently, it is not possible to ascertain the true number of people living at different stages of the condition in India based on clinical records alone. Efforts have been conducted to improve diagnosis rates in a country with such educational and linguistic diversity. Culturally appropriate tools such as the Indian Council of Medical Research-Neurocognitive Toolbox (ICMR-NCTB) (Iyer et al., 2020; Verma et al., 2021) have been developed and validated to aid in the establishment of accurate dementia diagnosis and prevalence estimates.

Care has a cost, and care needs will inevitably grow at an individual level as dementia progresses and impairment increases. For example, it has previously been estimated that care (e.g., formal care, informal care and medical costs) costs 45,600 Indian Rupees per year for a person with mild dementia (in an urban setting), whilst it costs over 2,02,450 Indian Rupees a year for a person with severe dementia (Rao & Bharath, 2013). The increased cost of care between severity was driven in part by increased residential care. The authors also report a lower cost of care in rural settings, which could be attributed to a lack of access to institutional care outside of metropolitan areas (Alzheimer's and Related Disorders Society of India, 2010). Irrespective, informal care typically is the largest cost for people with dementia in low- and middle- income settings, more so than social sector costs and direct medical costs (Wimo et al., 2018). There is much needed to understand the associated costs and impact of dementia in India to facilitate appropriate resource allocation and planning of service delivery.

This study aimed to generate and achieve expert consensus on:

- 1) The dementia prevalence for those aged 60 years and older in India, split by sex and age.
- 2) The percentage of people living with mild, moderate and severe dementia in India.

## **Methods**

We adopted a methodology similar to the one adopted in a global prevalence of dementia study (Ferri et al., 2005). Unlike a traditional Delphi study, we aimed to achieve consensus on estimates of dementia prevalence and dementia severity split through a series of unbound integer responses (i.e., expert provides numeric estimates, rather than being asked to rate their agreement or confidence on a specific estimates).

Eight experts were selected by a senior Indian neurologist and co-author (XX) on the basis of their experience and knowledge of the dementia field in the country. There is no consensus on the number of experts required for a Delphi study. For example, one study recommended that ten to fifteen experts is considered sufficient, though it should be acknowledged that smaller numbers do minimise potential logistical issues of running the Delphi technique (see Nashir et al., 2015). As per existing guidance (Adler & Ziglio, 1996; Skulmoski et al., 2007), experts were required to:

- (i) have the appropriate knowledge and experience,
- (ii) be willing to participate,
- (iii) have sufficient time (to participate) and,
- (iv) possess effective communication skills.

We prioritised experts with experience of research associated with estimating dementia prevalence within India. These experts were selected through snowballing techniques and utilising existing networks. Those selected as experts were not involved in the design, set up, or analysis involved in the Delphi study.

Each expert was sent an email from co-author (XX) describing the Delphi process and how their information/contributions would be treated if they decided to participate. All experts were informed that their individual responses would remain anonymised, though their involvement in the process would be acknowledged in any write-up (if they so wished).

In the first round, experts were provided a summary of evidence of prevalence data derived from the country (as identified in a previous publication; Farina et al., 2020). These prevalence data consisted of methodological details (setting, sample size, one-phase or two-phase survey, implementation of two-phase design, response rates for both phases, diagnostic criteria) and we summarised specific prevalence estimates from every study. We made no judgement about the quality or risk of bias of each study. The experts were explicitly informed that they do not need to use the prevalence data compiled. In a pre-designed form on REDCap (Harris et al.,

2009, 2019) hosted at the London School of Economics and Political Science, participants were asked to estimate the prevalence of dementia based upon their knowledge and/or using the information provided. These fields were split by age and sex. A minimum age of 60 years old was used to reflect that the vast majority of dementia cases occur in old age, whilst also aligning with many epidemiological studies that estimate dementia prevalence within India. Age categories were based on the available population demography from India (i.e., 60-64, 65-69, 70-74, 75-79, 80+).

Experts were also asked to provide an estimate of the percentage of people living with dementia in India, categorised by severity. We followed The Longitudinal Ageing Study in India (LASI) protocol that calculated the percentage in the lowest 10th percentile of the composite cognition score, as this was identified “as a proxy measure of poor cognitive functioning” (International Institute for Population Sciences et al., 2020, pp.374). From the composite cognition score we excluded serial 7s and backward counting from the original arithmetic domain due to missing values and to avoid our score being more relevant to measure education attainment than cognitive functioning respectively. After this, we operationalised the definition of severity of impairment by combining 10th percentile of the composite cognition score with Activities of Daily Living [ADL] (difficulty in walking across the room, bathing, eating, dressing, getting in or out of bed, or toilet use) as measured in the LASI dataset. This is in line with previous methodology (Comas-Herrera et al., 2007). The experts were provided the following guidance on how we defined severity using the LASI dataset:

- 1) Mild impairment: 10th percentile of composite cognition score and 1 ADL
- 2) Moderate impairment: 10th percentile of composite cognition score and 2 ADL
- 3) Severe impairment: 10th percentile of composite cognition score and 3 plus ADL

Even if experts were unsure, they were asked to provide their best estimate. Experts had the opportunity to add comments to support why they believed their estimates to be accurate. Alongside each set of responses, the experts were asked to self-report their confidence in their estimate (see Adler & Ziglio, 1996). Once all responses were received, data were compiled and summarised.

In the second round, experts were provided with a summary of evidence from the first round and were asked to estimate the prevalence in fields that did not reach consensus (see definition below). In this round, experts were provided information about the average responses (mean), and how their revised scores compared to the average response in the previous round. In

addition, a summary of anonymised comments alongside confidence ratings were provided. Experts were informed that they did not have to change their estimates if they did not want to.

It was decided that the Delphi process would stop once a) consensus was achieved across all fields, b) we had less than 70% response rates following a given round (thus affecting the study's validity), c) following three rounds irrespective of whether consensus had been achieved across all fields.

In this study we did not do a third round because the response rate dropped below 70% in the second round.

#### *Achieving consensus*

Consensus was determined by using an established *a priori* statistical criteria used in previous Delphi studies (von der Gracht, 2012). For consensus to be achieved in a given field following each round, the experts pooled estimates require to: a) achieve an Interquartile range (IQR) < 1, or b) the co-efficient of variation (CoV)  $\leq 0.5$ . The choice to utilise both criteria ensured that consensus could be achieved even in situations where estimate values were high, or where there were a small number of extreme outliers. The mean estimate for each field were extracted and used as the value that represented consensus to accommodate inevitable variations in estimates.

#### *Analysis and reporting*

Findings were summarised narratively for each round, with fields being grouped together based on theme (i.e., prevalence split by age, dementia severity) with an emphasis on fields that achieved a consensus. Fields that did not achieve consensus were also discussed, and the implications of this were considered. For each round and response, a mean, median, standard deviation, CoV and IQR were calculated and reported. Summary statistics of experts' confidence in their estimates were also reported in round one.

#### *Missing data*

During each round, we followed-up with experts if they did not respond or there were any gaps in the responses provided. If experts did not respond after three emails, then they were classed as drop-out.

#### *Maintaining anonymity*



The researcher (XX) who developed the forms and analysed data, did not have access to the names of the experts who responded to each round. A second researcher, who was not involved in the analysis process (XX), was responsible for liaising with the experts via email. All emails were sent out individually or blinded to other experts. In instances where the expert provided comments which could have revealed their identity, minor changes were made to maintain anonymity (e.g., "...based on our previous research (citation)" to "...based on previous research (citation)").

### *Ethical approvals*

All experts provided informed consent to share their expertise as part of the Delphi process. As this manuscript involves expert opinion rather than research on human subjects, we did not obtain ethical approval.

## **Results**

### **Round 1: Overview**

Of the 13 experts initially invited, eight expert responses were received. See Supplementary Table 1 for a summary of expertise.

#### **Prevalence split by sex and age**

Individual experts' estimates ranged from 0.5% (prevalence of males aged 60-64) to 27% (prevalence of males aged 80 years and over). Irrespective of sex, the average estimate increased by age category. Variability of estimates remained quite high, with no field meeting the *a priori* threshold of consensus either based on the IQR or CoV. See Table 1.

#### **Total prevalence**

Overall based on round 1 estimates, dementia prevalence was 4.9% for those aged 60 years and over in India, with a prevalence of 5.0% for males and 4.8% for females. Five of the experts were confident with their responses (60-79% confident of being right), two were very confident (80%-99% confident of being right), whilst one expert was less so (40-59% confident of being right).

#### **Severity**

A single expert's data entry had to be excluded because of a typographical error, leaving seven experts providing estimates on severity. Experts on average estimated that the largest proportion of people will be living with mild dementia (42.9%), followed by moderate (31.9%) and severe (25.2%). The mild and moderate severity estimates met the CoV threshold for consensus, though the severe estimate did not. See Table 2 for further details. Three experts were very confident with their estimates (80%-99% confident of being right), two experts were confident (60-79% confident of being right), one expert was unsure (40-59% confident of being right), and a single expert did not provide a confidence rating.

## **Round 2: Overview**

Between rounds three experts dropped out (non-response), thus leaving five experts for round two. Of the three experts that dropped out between rounds, two were estimating higher than average prevalence in first round (i.e., > 5.0% for total male prevalence and 4.8% for total female prevalence, respectively).

### **Prevalence split by sex and age**

In round two, consensus was achieved across experts on male prevalence split by age, ranging from 0.9% in the 60–64-year-old category to 9.4% in the 80 years and older category. For female estimates all estimates reached consensus apart from the 60–64-year-old group. Consistently for each age category, females were estimated to have an equal or higher dementia prevalence than males. Table 3 presents full details about average estimates and variability.

### **Total prevalence**

Following the second round, the average estimated prevalence for males was 2.4% (95% CI = 1.9 to 3.0), and 3.1% (95% CI = 2.0 to 4.2) for females. The total estimate of dementia prevalence in the over 60s in India was 2.8% (95% CI = 1.9 to 3.6).

### **Severity**

As consensus was achieved for estimates of mild and moderate severity in the first round, the emphasis was on determining whether consensus could be achieved for estimating the percentage of people currently living with severe dementia. In the second round, estimates of

dementia severity remained largely unchanged, with the mean estimate of severe dementia was 21.3% (SD=14.4, Mdn = 17.5, IQR = 18.8, CoV = 0.7). Consensus for this field was not achieved.

## **Discussion**

There are considerable gaps in evidence with respect to dementia prevalence in India. Among available evidence, there are limited data to allow prevalence estimates at a granular level. This study presented dementia prevalence estimates split by age and sex, but also the proportion of older adults potentially living at different stages of the condition in India. This was achieved using the Delphi process, through generating consensus among experts with substantial experience in epidemiology, neurology and public health in India. These prevalence estimates can aid in understanding the economic impact of dementia to society and further planning for appropriate resource allocation towards care services for dementia in India.

Our consensus estimates generated a dementia prevalence of 2.8% (95% CI = 1.9 to 3.6) for those aged 60 years and above in India. This would equate to approximately 3.9 million people living with dementia in India, when based on population projections for older adults living in India in 2021 (National Commission on Population, 2020). One systematic review estimated the dementia prevalence to be 2% in India<sup>1</sup> (Choudhary et al., 2021). Despite high quality synthesis, we should be cautious not to assume that such estimates are definitive. For example a recent logistic model for dementia status from LASI, which reported a prevalence of 7.4% for those aged 60 years and above in India (N = 28,949)(Lee et al., 2023). Such variability could be attributed to differences in demographics, region studied, methodology and diagnostic criteria employed (Lee et al., 2023; Ravindranath & Sundarakumar, 2021). The benefit of the Delphi process is that it allows for experts to factor this variability and recognise methodology that might underestimate the true prevalence of dementia.

Furthermore, this Delphi study sought to determine age and sex-specific prevalence estimates for dementia in India. Consensus was not achieved for the prevalence estimate of females aged 60-64. However, in line with previous studies (GBD 2019 Dementia Forecasting Collaborators, 2022; Prince et al., 2013), our consensus estimates demonstrated higher prevalence of dementia

---

<sup>1</sup> The meta-analysis was not limited by age and included data from studies that estimated prevalence in those aged  $\geq 50$  years.

in women than in men (3.1% vs 2.4). This may be partially attributed to longer life expectancy in women (Mielke, 2018), or poorer performance on cognitive tasks introduced through lower education attainment in women (Farrell et al., 2020; Zhang, 2006). Furthermore, in comparison to the GBD study (Singh et al., 2021), our consensus estimates were generally within the 95% confidence intervals reported, with only three estimates outside: males between 60 to 64 years of age, males between 65 to 69 years of age, and females aged 80 plus. Therefore, this strengthens the likelihood of our findings being tenable.

In addition to determining dementia prevalence split by age and sex, we also attempted to gain expert consensus on the proportion of those living at different stages of dementia in India. Severity of dementia is known to have a profound impact on estimating costs of care, with a prior study identifying an increase in dementia severity to contribute to a reduction in the proportion of medical costs, but an increase in care-related costs in India (Rao and Bharat, 2013). While no large epidemiological studies have examined dementia severity in the Indian population, a few studies conducted in specific regions of the country offer some insights (Banerjee et al., 2017; Rodriguez et al., 2008). For example, out of the 103 people detected with dementia in Kolkata City, 21.4% had mild, 25.2% had moderate and 53.4% had severe dementia (Banerjee et al., 2017). In our study, the experts estimated that 42.5% of people living with dementia had mild dementia (approximately 1.7 million older adults in India), and 31.9% had moderate dementia (approximately 1.2 million older adults in India). Consensus was not achieved for the percentage of persons living with severe dementia. The inability to achieve consensus in this category, may also be due to absence of relevant data or the use of different datasets by individual experts to estimate severity. For example, one expert broadly based their severity estimates on one study (Saldanha et al, 2010). Overall, understanding how the consensus estimates on dementia severity in India compare to other data is difficult, not least because there are no equivalent data existing in India. In the UK, a similar Delphi consensus processes estimated that from people with late-onset dementia, 55.4% have mild dementia and 32.1% have moderate dementia (Prince et al., 2014). In the US population-based Framingham Heart Study, researchers estimated that 50.4% of Alzheimer's disease participants had mild severity and 30.3% had moderate severity (Yuan et al., 2021). There is a need for further research in India to determine severity profile of dementia and allow for experts to more accurately estimate severity.

One of the key strengths of this study is that it is the first to use the Delphi method to estimate dementia prevalence and severity in India. The range of experts that have contributed to this study (including prominent neurologists, experienced in epidemiological research and diagnostic tools in India) has provided some validity to our consensus estimates, which have added to the current evidence base on dementia prevalence and severity. We utilized *a priori* criteria to define how consensus was achieved, and when the Delphi process should be stopped. However, it should also be acknowledged that there is no current gold standard threshold, and that applying different definitions of consensus could yield different results.

Other limitations are also recognized. The Delphi process did not continue to the third round due the dropout rates between round 1 and 2 exceeding 30%, thus potentially affecting the validity of the findings. Experts that dropped out between rounds tended to have higher prevalence estimates in the first round, and hence it is possible that they did not feel comfortable with altering their score to align with others. Another caveat of the process is that the consensus estimates are based completely on expert opinion, although efforts were made to minimise potential bias. As experts draw conclusions largely based on the existing evidence, possible conclusions are considerably dependent on this limited and variable evidence. The process is also susceptible to cognitive biases such as the bandwagon effect (in which people will change their responses just so they align with others, irrespective of their own belief) and belief perseverance (in which people will not change their beliefs even in light of new information) (Winkler & Moser, 2016). The extent to which the experts were given direction to formulate their estimates, may have also influenced their estimates. For example, we did not provide guidance on the diagnostic criteria that should be used to estimate prevalence, or explicitly define dementia. It is unclear the extent to which this lack of guidance may have shaped experts' decisions regarding the estimates generated. All experts had backgrounds in medicine, thus potentially minimising the adoption of non-clinical definitions and diagnostic criteria of dementia. As with any Delphi process, the choice of experts is likely to influence the conclusions made, as their input will be based on their training and experiences.

In conclusion, our findings provide detailed information regarding the number of people living with dementia in India and also novel insights into how many people are potentially living at each stage of the condition. Such data can be utilized to estimate the economic impact of dementia and also allow for better allocation of resources towards dementia care in India until further primary evidence is generated.

## **Acknowledgements**

We would like to thank the following experts for contributing towards this Delphi study: **Dr Tapas Kumar Banerjee**, Medical Director & Chief Consultant Neurologist, National Neurosciences Centre Calcutta, Kolkata; **Dr K Chandra Sekhar**, Managing Director & Senior Consultant, Geriatric psychiatry and Neuropsychiatry, Asha Hospital, Hyderabad and President-ARDSI Hyderabad Deccan chapter; **Dr Robert Mathew**, Professor of Neurology Sree Mookambika Institute of Medical Sciences, Kanyakumari District, Tamil Nadu; **Dr Yogeshwar Kalkonde**, Sangwari, Surguja, Chhattisgarh; **Dr Amitabha Ghosh**, Director and Head, Department of Neurology, Apollo Multispeciality Hospitals, Kolkata; **Dr Atanu Biswas**, Professor, Department of Neurology, Institute of Post Graduate Medical Education & Research and Bangur Institute of Neurosciences, Kolkata; **Prof Lalit Dandona**, Public Health Foundation of India, Gurugram, India; Indian Council of Medical Research, New Delhi, India; and Institute for Health Metrics and Evaluation, University of Washington, Seattle, USA; **Dr Ramshekhar Menon**, Department of Neurology, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Thiruvananthapuram, India. We would like to acknowledge that the experts did not review or approve the final manuscript prior to publication. We would like to thank **Dr Paulino Font-Gilabert** (King's College London) for his input and contribution to the study.

## **Funding**

This work was supported by the UK Research and Innovation's Global Challenges Research Fund (ES/P010938/1). The funder has not influenced the design, outcome or interpretation of the study.

## References

- Adler, M., & Ziglio, E. (1996). *Gazing Into the Oracle: The Delphi Method and Its Application to Social Policy and Public Health*. Jessica Kingsley Publishers.
- Alzheimer's and Related Disorders Society of India. (2010). *The Dementia India Report 2010: Prevalence, impact, costs and services for dementia*. Alzheimer's and Related Disorders Society of India.
- Banerjee, T. K., Dutta, S., Das, S., Ghosal, M., Ray, B. K., Biswas, A., Hazra, A., Chaudhuri, A., Paul, N., & Das, S. K. (2017). Epidemiology of dementia and its burden in the city of Kolkata, India. *International Journal of Geriatric Psychiatry*, 32(6), 605–614. <https://doi.org/10.1002/gps.4499>
- Chandra, V., Ganguli, M., Pandav, R., Johnston, J., Belle, S., & DeKosky, S. T. (1998). Prevalence of Alzheimer's disease and other dementias in rural India: The Indo-US study. *Neurology*, 51(4), 1000–1008. <https://doi.org/10.1212/wnl.51.4.1000>
- Choudhary, A., Ranjan, J. K., & Asthana, H. S. (2021). Prevalence of dementia in India: A systematic review and meta-analysis. *Indian Journal of Public Health*, 65(2), 152. [https://doi.org/10.4103/ijph.IJPH\\_1042\\_20](https://doi.org/10.4103/ijph.IJPH_1042_20)
- Comas-Herrera, A., Wittenberg, R., Pickard, L., & Knapp, M. (2007). Cognitive impairment in older people: Future demand for long-term care services and the associated costs. *International Journal of Geriatric Psychiatry*, 22(10), 1037–1045. <https://doi.org/10.1002/gps.1830>
- Das, S. K., Biswas, A., Roy, J., Bose, P., Roy, T., Banerjee, T. K., Mukherjee, C., Raut, D. K., Chowdhury, A., & Hazra, A. (2008). Prevalence of major neurological disorders among geriatric population in the metropolitan city of Kolkata. *The Journal of the Association of Physicians of India*, 56, 175–181.

- Das, S. K., Biswas, A., Roy, T., Banerjee, T. K., Mukherjee, C. S., Raut, D. K., & Chaudhuri, A. (2006). A random sample survey for prevalence of major neurological disorders in Kolkata. *The Indian Journal of Medical Research*, *124*(2), 163–172.
- Das, S. K., Pal, S., & Ghosal, M. K. (2012). Dementia: Indian scenario. *Neurology India*, *60*(6), 618. <https://doi.org/10.4103/0028-3886.105197>
- Farina, N., Ibdidris, A., Alladi, S., Comas-Herrera, A., Albanese, E., Docrat, S., Ferri, C. P., Freeman, E., Govia, I., Jacobs, R., Astudillo-Garcia, C. I., Musyimi, C., Sani, T. P., Schneider, M., Theresia, I., Turana, Y., Knapp, M., & and, S. B. (2020). A systematic review and meta-analysis of dementia prevalence in seven developing countries: A STRiDE project. *Global Public Health*, 1–16. <https://doi.org/10.1080/17441692.2020.1792527>
- Farrell, M. T., Kobayashi, L. C., Montana, L., Wagner, R. G., Demeyere, N., & Berkman, L. (2020). Disparity in Educational Attainment Partially Explains Cognitive Gender Differences in Older Rural South Africans. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, *75*(7), e161–e173. <https://doi.org/10.1093/geronb/gbaa035>
- Ferri, C. P., Prince, M., Brayne, C., Brodaty, H., Fratiglioni, L., Ganguli, M., Hall, K., Hasegawa, K., Hendrie, H., Huang, Y., Jorm, A., Mathers, C., Menezes, P. R., Rimmer, E., & Scazufca, M. (2005). Global prevalence of dementia: A Delphi consensus study. *The Lancet*, *366*(9503), 2112–2117. [https://doi.org/10.1016/S0140-6736\(05\)67889-0](https://doi.org/10.1016/S0140-6736(05)67889-0)
- GBD 2019 Dementia Forecasting Collaborators. (2022). Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: An analysis for the Global Burden of Disease Study 2019. *The Lancet. Public Health*, *7*(2), e105. [https://doi.org/10.1016/S2468-2667\(21\)00249-8](https://doi.org/10.1016/S2468-2667(21)00249-8)



- Harris, P. A., Taylor, R., Minor, B. L., Elliott, V., Fernandez, M., O'Neal, L., McLeod, L., Delacqua, G., Delacqua, F., Kirby, J., Duda, S. N., & REDCap Consortium. (2019). The REDCap consortium: Building an international community of software platform partners. *Journal of Biomedical Informatics*, *95*, 103208. <https://doi.org/10.1016/j.jbi.2019.103208>
- Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics*, *42*(2), 377–381. <https://doi.org/10.1016/j.jbi.2008.08.010>
- International Institute for Population Sciences, National Programme for Health Care of Elderly, MoHFW, Harvard T. H. Chan School of Public Health, & University of Southern California. (2020). *Longitudinal Ageing Study in India (LASI) Wave 1, 2017-18, India Report*. International Institute for Population Sciences.
- Iyer, G. K., Paplikar, A., Alladi, S., Dutt, A., Sharma, M., Mekala, S., Kaul, S., Saroja, A. O., Divyaraj, G., Ellajosyula, R., Ghosh, A., Hooda, R., Justus, S., Kandukuri, R., Khan, A. B., Mathew, R., Mathuranath, P. S., Menon, R., Nandi, R., ... Consortium, I. N. T. B. (2020). Standardising Dementia Diagnosis Across Linguistic and Educational Diversity: Study Design of the Indian Council of Medical Research-Neurocognitive Tool Box (ICMR-NCTB). *Journal of the International Neuropsychological Society*, *26*(2), 172–186. <https://doi.org/10.1017/S1355617719001127>
- Lee, J., Meijer, E., Langa, K. M., Ganguli, M., Varghese, M., Banerjee, J., Khobragade, P., Angrisani, M., Kurup, R., Chakrabarti, S. S., Gambhir, I. S., Koul, P. A., Goswami, D., Talukdar, A., Mohanty, R. R., Yadati, R. S., Padmaja, M., Sankhe, L., Rajguru, C., ... Dey, A. B. (2023). Prevalence of dementia in India: National and state

estimates from a nationwide study. *Alzheimer's & Dementia*.

<https://doi.org/10.1002/alz.12928>

- Mathuranath, P. S., Cherian, P. J., Mathew, R., Kumar, S., George, A., Alexander, A.,  
Ranjith, N., & Sarma, P. S. (2010). Dementia in Kerala, South India: Prevalence and  
influence of age, education and gender. *International Journal of Geriatric Psychiatry*,  
25(3), 290–297. <https://doi.org/10.1002/gps.2338>
- Mielke, M. M. (2018). Sex and Gender Differences in Alzheimer's Disease Dementia. *The  
Psychiatric Times*, 35(11), 14–17.
- Nashir, I. M., Mustapha, R., & Yusoff, A. (2015). Delphi technique: Enhancing research in  
technical and vocational education. *Journal of Technical Education and Training*,  
7(2).
- National Commission on Population. (2020). *Population Projections for India and States  
2011-2036. Report of the Technical Group on Population Projections*. Ministry of  
Health and Family Welfare.  
[https://main.mohfw.gov.in/sites/default/files/Population%20Projection%20Report%202011-2036%20-%20upload\\_compressed\\_0.pdf](https://main.mohfw.gov.in/sites/default/files/Population%20Projection%20Report%202011-2036%20-%20upload_compressed_0.pdf)
- Nulkar, A., Paralikar, V., & Juvekar, S. (2019). Dementia in India – a call for action. *Journal  
of Global Health Reports*, 3, e2019078. <https://doi.org/10.29392/joghr.3.e2019078>
- Paddick, S.-M., Longdon, A. R., Kisoli, A., Dotchin, C., Gray, W. K., Dewhurst, F., Chaote,  
P., Kalaria, R., Jusabani, A. M., & Walker, R. (2013). Dementia prevalence estimates  
in sub-Saharan Africa: Comparison of two diagnostic criteria. *Global Health Action*,  
6, 10.3402/gha.v6i0.19646. <https://doi.org/10.3402/gha.v6i0.19646>
- Perianayagam, A., Bloom, D., Lee, J., Parasuraman, S., Sekher, T. V., Mohanty, S. K.,  
Chattopadhyay, A., Govil, D., Pedgaonkar, S., Gupta, S., Agarwal, A., Posture, A.,  
Weerman, A., & Pramanik, S. (2022). Cohort Profile: The Longitudinal Ageing Study

in India (LASI). *International Journal of Epidemiology*, dyab266.

<https://doi.org/10.1093/ije/dyab266>

Poddar, K., Kant, S., Singh, A., & Singh, T. B. (2011). An epidemiological study of dementia among the habitants of eastern Uttar Pradesh, India. *Annals of Indian Academy of Neurology*, 14(3), 164–168. <https://doi.org/10.4103/0972-2327.85874>

Prince, M., Bryce, R., Albanese, E., Wimo, A., Ribeiro, W., & Ferri, C. P. (2013). The global prevalence of dementia: A systematic review and metaanalysis. *Alzheimer's & Dementia*, 9(1), 63-75.e2. <https://doi.org/10.1016/j.jalz.2012.11.007>

Prince, M., Knapp, M., Guerchet, M., McCrone, P., Prina, M., Comas-Herrera, A., Wittenberg, R., Adelaja, B., Hu, B., King, D., Rehill, A., & Salimkumar, D. (2014). *Dementia UK: Update* (Second). Alzheimer's Society.

Raina, S. K., Chander, V., & Bhardwaj, A. (2016). Dementia in a tribal landlocked elderly population at high altitude: What explains the lower prevalence? *Journal of Neurosciences in Rural Practice*, 7(3), 419–422. <https://doi.org/10.4103/0976-3147.182775>

Raina, S. K., Raina, S., Chander, V., Grover, A., Singh, S., & Bhardwaj, A. (2014). Is dementia differentially distributed? A study on the prevalence of dementia in migrant, urban, rural, and tribal elderly population of himalayan region in northern India. *North American Journal of Medical Sciences*, 6(4), 172–177.

<https://doi.org/10.4103/1947-2714.131243>

Raina, S., Raina, S., Razdan, S., & Pandita, K. (2008). Prevalence of dementia among Kashmiri migrants. *Annals of Indian Academy of Neurology*, 11(2), 106.

<https://doi.org/10.4103/0972-2327.41878>

Rajkumar, S., Kumar, S., & Thara, R. (1997). Prevalence of dementia in a rural setting: A report from India. *International Journal of Geriatric Psychiatry*, 12(7), 702–707.

- Rao, G. N., & Bharath, S. (2013). Cost of dementia care in India: Delusion or reality? *Indian Journal of Public Health, 57*(2), 71. <https://doi.org/10.4103/0019-557X.114986>
- Ravindranath, V., & Sundarakumar, J. S. (2021). Changing demography and the challenge of dementia in India. *Nature Reviews Neurology, 17*(12), Article 12. <https://doi.org/10.1038/s41582-021-00565-x>
- Razdan, S., Kaul, R. L., Motta, A., Kaul, S., & Bhatt, R. K. (1994). Prevalence and Pattern of Major Neurological Disorders in Rural Kashmir (India) in 1986. *Neuroepidemiology, 13*(3), 113–119. <https://doi.org/10.1159/000110368>
- Rodriguez, J. J. L., Ferri, C. P., Acosta, D., Guerra, M., Huang, Y., Jacob, K., Krishnamoorthy, E., Salas, A., Sosa, A. L., Acosta, I., Dewey, M. E., Gaona, C., Jotheeswaran, A., Li, S., Rodriguez, D., Rodriguez, G., Kumar, P. S., Valhuerdi, A., Prince, M., & 10/66 Dementia Research Group. (2008). Prevalence of dementia in Latin America, India, and China: A population-based cross-sectional survey. *The Lancet, 372*(9637), 464–474. [https://doi.org/10.1016/S0140-6736\(08\)61002-8](https://doi.org/10.1016/S0140-6736(08)61002-8)
- Saldanha, D., Mani, M. R., Srivastava, K., Goyal, S., & Bhattacharya, D. (2010). An epidemiological study of dementia under the aegis of mental health program, Maharashtra, Pune chapter. *Indian Journal of Psychiatry, 52*(2), 131–139. <https://doi.org/10.4103/0019-5545.64588>
- Seby, K., Chaudhury, S., & Chakraborty, R. (2011). Prevalence of psychiatric and physical morbidity in an urban geriatric population. *Indian Journal of Psychiatry, 53*(2), 121.
- Shaji, S., Bose, S., & Verghese, A. (2005). Prevalence of dementia in an urban population in Kerala, India. *British Journal of Psychiatry, 186*(2), 136–140. <https://doi.org/10.1192/bjp.186.2.136>
- Singh, G., Sharma, M., Kumar, G. A., Rao, N. G., Prasad, K., Mathur, P., Pandian, J. D., Steinmetz, J. D., Biswas, A., Pal, P. K., Prakash, S., Sylaja, P. N., Nichols, E., Dua,

- T., Kaur, H., Alladi, S., Agarwal, V., Aggarwal, S., Ambekar, A., ... Dandona, L. (2021). The burden of neurological disorders across the states of India: The Global Burden of Disease Study 1990–2019. *The Lancet Global Health*, 9(8), e1129–e1144. [https://doi.org/10.1016/S2214-109X\(21\)00164-9](https://doi.org/10.1016/S2214-109X(21)00164-9)
- Skulmoski, G. J., Hartman, F. T., & Krahn, J. (2007). The Delphi Method for Graduate Research. *Journal of Information Technology Education: Research*, 6, 001–021.
- Vas, C. J., Pinto, C., Panikker, D., Noronha, S., Deshpande, N., Kulkarni, L., & Sachdeva, S. (2001). Prevalence of Dementia in an Urban Indian Population. *International Psychogeriatrics*, 13(4), 439–450. <https://doi.org/10.1017/S1041610201007852>
- Verma, M., Tripathi, M., Nehra, A., Paplikar, A., Varghese, F., Alladi, S., Narayanan, J., Dhaliwal, R. S., Sharma, M., Saroja, A. O., Arshad, F., Divyaraj, G., Ghosh, A., Manae, T. S., Mekala, S., Menon, R. N., Hooda, R., Iyer, G. K., Sunitha, J., ... Vishwanath, Y. (2021). Validation of ICMR Neurocognitive Toolbox for Dementia in the Linguistically Diverse Context of India. *Frontiers in Neurology*, 12, 661269. <https://doi.org/10.3389/fneur.2021.661269>
- von der Gracht, H. (2012). Consensus measurement in Delphi studies Review and implications for future quality assurance. *Technological Forecasting and Social Change*, 79, 1525–1536. <https://doi.org/10.1016/j.techfore.2012.04.013>
- Wimo, A., Gauthier, S., Prince, M., & on behalf of ADI's Medical Scientific Advisory. (2018). *Global estimates of informal care*. Alzheimer's Disease International and Karolinska Institutet.
- Winkler, J., & Moser, R. (2016). Biases in future-oriented Delphi studies: A cognitive perspective. *Technological Forecasting and Social Change*, 105, 63–76. <https://doi.org/10.1016/j.techfore.2016.01.021>

- Yuan, J., Maserejian, N., Liu, Y., Devine, S., Gillis, C., Massaro, J., & Au, R. (2021). Severity Distribution of Alzheimer's Disease Dementia and Mild Cognitive Impairment in the Framingham Heart Study. *Journal of Alzheimer's Disease*, 79(2), 807–817. <https://doi.org/10.3233/JAD-200786>
- Zhang, Z. (2006). Gender Differentials in Cognitive Impairment and Decline of the Oldest Old in China. *The Journals of Gerontology: Series B*, 61(2), S107–S115. <https://doi.org/10.1093/geronb/61.2.S107>

Table 1. Average estimate of dementia prevalence in India split by age and sex after the first round with experts (n=8).

Age	Sex	Mean	SD	Mdn	IQR	CoV	Meet <i>a priori</i> consensus
60-64	Male	2.6	3.2	1.5	2.0	1.2	No
	Female	2.0	1.5	1.6	3.1	0.8	No
65-69	Male	3.9	4.8	2.3	3.9	1.2	No
	Female	3.8	3.6	2.5	3.8	1.0	No
70-74	Male	6.0	6.6	3.3	4.5	1.1	No
	Female	5.5	4.8	3.8	6.5	0.9	No
75-79	Male	7.7	6.2	4.8	5.6	0.8	No
	Female	8.0	6.2	5.2	6.9	0.8	No
80+	Male	11.9	7.8	9.6	10.5	0.7	No
	Female	12.2	7.1	10.2	10.7	0.6	No

Table 2. Estimate of the percentage of people living with dementia in India split by severity (n=7).

	Mean	SD	Mdn	IQR	CoV	Meet <i>a</i> <i>priori</i> consensus
Mild	42.9	21.5	40	45.0	0.5	Yes
Moderate	31.9	10.8	30	17.5	0.3	Yes
Severe	25.2	15.6	25	27.5	0.6	No



Table 3. Average estimate of dementia prevalence (%) in India split by age and sex after the second round with experts (n=5).

Age	Sex	Mean	SD	Mdn	IQR	CoV	Meet <i>a priori</i> consensus
60-64	Male	0.9	0.5	0.6	0.5	0.6	Yes
	Female	0.9	0.5	0.8	1.1	0.6	No
65-69	Male	1.4	0.4	1.2	0.6	0.3	Yes
	Female	1.5	0.4	1.5	0.9	0.3	Yes
70-74	Male	2.1	0.3	2.0	0.1	0.1	Yes
	Female	3.0	0.4	3.2	0.8	0.1	Yes
75-79	Male	4.3	0.5	4.0	0.6	0.1	Yes
	Female	5.2	1.3	5.0	1.4	0.2	Yes
80+	Male	9.4	1.5	10.0	3.0	0.2	Yes
	Female	12.3	4.7	11.0	5.0	0.4	Yes

**Supplementary Table 1. Brief description of the experts that participated in the Delphi process.**

<b>Expert #</b>	<b>Expertise</b>
<b>1</b>	Neurologist, been a part of dementia epidemiological studies conducted in the Indian context
<b>2</b>	Neurologist, been a part of dementia epidemiological studies conducted in the Indian context
<b>3</b>	Physician and Research Professor, been a part of national and international level research on burden of neurological disorders.
<b>4</b>	Leads a chapter of an Alzheimer's related NGO in India and is a neurologist.
<b>5</b>	Neurologist, has aided in a study validating culturally appropriate diagnostic tools for dementia/MCI
<b>6</b>	Neurologist, has aided in a study validating culturally appropriate diagnostic tools for dementia/MCI
<b>7</b>	Public health researcher and a neurologist
<b>8</b>	Geriatric Psychiatrist and leads a chapter of Alzheimer's related NGO in India