



PEARL

DERMOPHIS GLANDULOSUS (Glandular Caecilian) DIET

Puschendorf, R; Chaves, G

Published in:

The Herpetological Bulletin

Publication date:

2016

Link:

[Link to publication in PEARL](#)

Citation for published version (APA):

Puschendorf, R., & Chaves, G. (2016). DERMOPHIS GLANDULOSUS (Glandular Caecilian) DIET. *The Herpetological Bulletin*, 126(0), 31-31.

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

DERMOPHIS GLANDULOSUS (Glandular Caecilian) DIET. As with most caecilians we know little about the biology of *Dermophis glandulosus*, other than its taxonomic status and its distribution from Costa Rica to northern Colombia (Savage & Wake 2001). This species seems to be one of the more rare caecilians for this region with just 10 specimens deposited in the collection of the Museum of Zoology at the University of Costa Rica. Herein we describe the stomach contents of a specimen found dead in the field on the 9 October 2005 near Rio J aris, Canton de Mora, Costa Rica (9.9043 N, 84.3012W; 550 m asl; in tropical humid forest).

The specimen was in good condition when found, however, cause of death could not be determined. Upon examination of its stomach contents we discovered a well-developed egg of an *Anolis* sp. It was identified as an *Anolis* egg as the embryo had well developed lamellae on all of its fingers (Losos, 2009); nevertheless it was impossible to identify the species as the body scales were not yet fully developed. The embryo presented 14 lamellae under the forth toe, which is common for most medium sized anoles in the region (Savage, 2002). The most abundant species in this area are *A. cupreus* and *A. intermedius*. Because *Anolis* eggs are commonly laid above ground or just under a

rock or log (Losos, 2009), this record suggests this species could be actively foraging on the forest floor and not just underground as it is thought of for most caecilians.

REFERENCES

- Losos, J.B. (2009). *Lizards in the Evolutionary Tree: Ecology and Adaptive Radiation of Anoles*. University of California Press. Berkeley, Los Angeles, London. Pp 507.
- Savage, J. (2002). *Amphibian and Reptiles of Costa Rica: A Herpetofauna between two Continents, between two Seas*. The University of Chicago Press. Chicago and London. Pp 934.
- Savage, J.M. & Wake, M.H. A. (2001) Reevaluation of the Status of Taxa of Central American Caecilians (Amphibia: Gymnophiona), with Comments on Their Origin and Evolution. *Copeia* 2001: 52-64.

Submitted by: ROBERT PUSCHENDORF, School of Biological Sciences, Plymouth University, Drake Circus, Plymouth, Devon, PL4 8AA, UK. robert.puschendorf@gmail.com, GERARDO CHAVES, Escuela de Biolog a, Universidad de Costa Rica, San Jos e, Costa Rica. Cachi13@gmail.com.