



PEARL

### Chemical analysis of flotsam ambergris

Rowland, Steven J.; Sutton, Paul A.; Lewis, C. Anthony; Knowles, Timothy; Wilde, Michael J.; Alves, Filipe; Clough, Robert

**Published in:**

Natural Product Research

**DOI:**

[10.1080/14786419.2024.2361863](https://doi.org/10.1080/14786419.2024.2361863)

**Publication date:**

2024

**Document version:**

Publisher's PDF, also known as Version of record


**Link:**

[Link to publication in PEARL](#)

**Citation for published version (APA):**

Rowland, S. J., Sutton, P. A., Lewis, C. A., Knowles, T., Wilde, M. J., Alves, F., & Clough, R. (2024). Chemical analysis of flotsam ambergris. *Natural Product Research*. <https://doi.org/10.1080/14786419.2024.2361863>

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Wherever possible please cite the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



# Natural Product Research

Formerly Natural Product Letters

ISSN: (Print) (Online) Journal homepage: [www.tandfonline.com/journals/gnpl20](http://www.tandfonline.com/journals/gnpl20)

## Chemical analysis of flotsam ambergris

Steven J. Rowland, Paul A. Sutton, C. Anthony Lewis, Timothy Knowles, Michael J. Wilde, Filipe Alves & Robert Clough

To cite this article: Steven J. Rowland, Paul A. Sutton, C. Anthony Lewis, Timothy Knowles, Michael J. Wilde, Filipe Alves & Robert Clough (01 Jul 2024): Chemical analysis of flotsam ambergris, Natural Product Research, DOI: [10.1080/14786419.2024.2361863](https://doi.org/10.1080/14786419.2024.2361863)

To link to this article: <https://doi.org/10.1080/14786419.2024.2361863>



© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



[View supplementary material](#)



Published online: 01 Jul 2024.



[Submit your article to this journal](#)



Article views: 958



[View related articles](#)



[View Crossmark data](#)



Citing articles: 1 [View citing articles](#)

## Chemical analysis of flotsam ambergris

Steven J. Rowland<sup>a</sup> , Paul A. Sutton<sup>a</sup> , C. Anthony Lewis<sup>a</sup> , Timothy Knowles<sup>b</sup> , Michael J. Wilde<sup>a</sup> , Filipe Alves<sup>c,d</sup>  and Robert Clough<sup>a</sup> 

<sup>a</sup>Biogeochemistry Research Centre, University of Plymouth, Drake Circus, Plymouth, UK; <sup>b</sup>Bristol Radiocarbon Accelerator Mass Spectrometry Facility (BRAMS), University of Bristol, Bristol, UK; <sup>c</sup>Regional Agency for the Development of Research Technology and Innovation (ARDITI), MARE – Marine and Environmental Sciences Centre/ARNET – Aquatic Research Network, Madeira, Portugal; <sup>d</sup>Faculty of Life Sciences, University of Madeira, Madeira, Portugal

### ABSTRACT

The natural product ambergris is only found rarely on beaches, as jetsam. Even more scarce, or even absent, are accounts of flotsam ambergris. Here, we report the chemical analysis of a rare, large piece (>100kg) of flotsam found in the Atlantic in 2019. About 95% of subsamples from the outside of the coprolith was soluble in dichloromethane. Of this, FTIR spectroscopy, APCI-MS and GC-MS indicated the presence of ambrein. Radiocarbon dating indicated that the sample was post 1950s in age. The  $^{13}\text{C}/^{12}\text{C}$  isotope ratio (−22.5 ‰) was typical of those reported to date for whale ‘body’ ambergris. Metals of ambergris have hardly been reported previously. The distribution found here for the flotsam, was dominated by copper and zinc, which is similar to that of several squid species. This is also consistent with the presence of squid beaks in the coprolith. Squid are a major prey species of sperm whales.


### ARTICLE HISTORY

Received 2 February 2024


Accepted 26 May 2024

### KEYWORDS

Ambergris; flotsam; *Physeter macrocephalus*; atlantic; ambrein; squid; copper



**CONTACT** Steven J. Rowland  [srowland@plymouth.ac.uk](mailto:srowland@plymouth.ac.uk)

 Supplemental data for this article can be accessed online at <https://doi.org/10.1080/14786419.2024.2361863>.

© 2024 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

## 1. Introduction

In 1954, in the journal *Nature*, Clarke reported a 'huge haul of ambergris' (Clarke 1954), which is a rare natural product of sperm whales *Physeter macrocephalus* (Cornon 1955, Clarke 2006, Mikhalev 2014). This 'body' ambergris was found inside a male whale in the Southern Ocean (Clarke 1954). Ambergris has also been rarely but consistently found as jetsam on beaches around the globe (Brito et al. 2016; Rowland et al. 2018, 2021; Wilde et al. 2020). Since it is found as jetsam, ambergris must logically also occur as flotsam, but so far as we are aware, no scientific reports of flotsam ambergris have been published.

Here, we studied the chemical composition of a floating boulder of ambergris (flotsam) retrieved from the Atlantic in 2019, which weighed over 100 kg (Figure S1). Flotsam ambergris is very rare, and this is an unusually large piece.

## 2. Results and discussion

Jetsam ambergris sometimes contains whole or fragments of squid beaks, which were abundant in the present flotsam sample (Figure S1). Scientific methods for the assignment of ambergris on the basis of ambrein (I) content, have been described in detail previously, including the use of radiocarbon dating (Rowland et al. 2019), Fourier transform infra-red spectroscopy (FTIR), gas chromatography-mass spectrometry (GC-MS) with and without derivatisation (Governo et al. 1977; Rowland and Sutton 2017),  $^1\text{H}$  and  $^{13}\text{C}$  nuclear magnetic resonance spectroscopy (Rowland et al. 2018), isotope ratio GC-MS (Rowland et al. 2021), atmospheric pressure chemical ionisation-MS (APCI-MS; Rowland et al. 2024) and DNA profiling (Macleod et al. 2020). In the present study of flotsam, several of the above methods were used to assign the material as ambergris.

For instance, radiocarbon dating (Table S1) indicated an age post-1950s (i.e. 'modern') with an  $\text{F}_{14\text{C}}$  value (1.0566) consistent with this (Rowland et al. 2019).

About  $95 \pm 3\%$  ( $n=3$ ) of each of three 40–60 mg subsamples was soluble in dichloromethane (DCM). This is typical of many samples of jetsam ambergris examined previously, which also had high contents of organic-soluble material (Rowland et al. 2019). The FTIR spectra of DCM extracts contained the typical features of spectra of ambergris solutions, dominated by absorptions due to the functional groups of ambrein (I). Thus, a broad transmittance at  $3372\text{cm}^{-1}$  was attributed to H-bonded hydroxyl O-H stretching. A weak transmittance at  $\sim 3067\text{cm}^{-1}$  was indicative of unsaturation and attributed to C-H stretch in an alkene. Transmittances at  $2925$  and  $2863\text{cm}^{-1}$  were attributed to C-H stretching in methyl and methylene groups and those at  $1461$  and  $1382\text{cm}^{-1}$  to the corresponding bending vibrations. Transmittances at  $1644\text{cm}^{-1}$  were attributed to C=C stretch and at  $887\text{cm}^{-1}$  to the =C-H out of plane bend (cf Governo et al. 1977; Rowland and Sutton 2017; Rowland et al. 2018).

Similarly, the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra (Figure S2) comprised resonances indicative of the distinctive features of ambrein, including those assigned to the alkenic H and C (cf Rowland et al. 2018; Rowland et al. 2018a). For example, resonances at 4 to about 5 ppm (Figure S2a) in the  $^1\text{H}$  NMR spectrum were assigned to alkenic protons. The two broad singlets occurring at 4.5 and 4.7 ppm were assigned to the two

methylenic protons (Rowland et al. 2018). Similarly, the triplet at about 5.2 ppm was assigned to the vinylic proton at 3' (Figure S2a). These characteristic resonances allow the presence of ambrein (and identity of the origin of the sample as ambergris) to be assigned. In addition, the resonances due to two methyl groups of ambrein were also assigned (Figure S2a). The presence of minor unknown constituents was indicated by other minor resonances in the region between 0.5 and ~2.2 ppm. These underlay the characteristic resonances of pure ambrein (Rowland et al. 2018). The  $^{13}\text{C}$  NMR spectrum of the whole dichloromethane extract of the flotsam ambergris re-dissolved in deuterated chloroform, is shown in Figure S2b. Resonances at ~100–150 ppm were assigned to alkenic carbon atoms by comparison with published spectra (Rowland et al. 2018). The resonances occurring in this region were each assigned to C6'', 4', 3' and the methylenic C6''=CH<sub>2</sub>. In addition, the resonance due to C2 was also assigned (Figure S2b).

GC-MS indicated an average composition of  $95 \pm 0.5\%$  ( $n=3$ ; Table S2) ambrein, measured and identified as the TMS ether (Figure S3; cf Rowland and Sutton 2017; Rowland et al. 2018; Rowland et al. 2018a). Thus, a small ion  $m/z$  485 was assigned to the  $\text{M}^+-\text{CH}_3$  ion of the TMS ether (Rowland and Sutton 2017), whilst a significant ion at  $m/z$  143 was attributed to a mono-unsaturated, C<sub>4</sub>H<sub>6</sub>-OTMS moiety (Rowland and Sutton 2017).

Positive ion APCI-MS (Figure S4; cf Rowland et al. 2024) also confirmed the presence of ambrein. The spectrum was characterised by a base peak ion,  $m/z$  411, shown previously by high resolution accurate mass APCI-MS to originate from the protonated molecular ion ( $m/z$  429;  $[\text{MH}]^+$ ) of the hindered alcohol ambrein (I) by water loss (Rowland et al. 2024).

The approximate  $^{13}\text{C}/^{12}\text{C}$  isotope ratio was  $-22.5\text{‰}$  (Table S1). (Repeat analyses established previously that the reproducibility of the latter method was  $\pm 0.4\text{‰}$ ; Rowland et al. 2019). For comparison, the stable isotopic compositions of the carbon in jetsam ambergris from both hemispheres examined previously had a mean value of  $-21.6 \pm 1.9\text{‰}$ ;  $n=26$ ; Rowland et al. 2019). The stable isotope ratios of blubber, skin, liver and muscle from seven northern hemisphere sperm whales from the Adriatic coast of Southern Italy was  $-18.5 \pm 1.2\text{‰}$  (Mazzariol et al. 2011).

ICP-MS revealed the relative and absolute concentrations of metals in the flotsam (Figure S5). Lederer (1950) postulated that the metals in ambergris might reflect the metal composition of the squid diet of the whale, particularly the squid haemolymph, in which copper complexed as haemocyanin, would be the major metal. Ohloff et al. (1977) suggested that copper porphyrins in ambergris would act as catalysts for photo- and auto-oxidation of ambrein; photo-oxidation experiments with 5, 10, 15, 20-tetraphenyl-21H, 23H-porphine copper (II), supported this (Rowland and Payne 2018). Few data for metal contents of ambergris have been published to date.

Copper was the major element detected in the flotsam ambergris studied herein ( $\sim 70\text{ }\mu\text{g g}^{-1}$  dried weight in the solid; Figure S5), but concentrations of zinc were similar ( $\sim 50\text{ }\mu\text{g g}^{-1}$ ), whilst iron and cadmium were present at appreciable, but somewhat lower, concentrations ( $\sim 20\text{ }\mu\text{g g}^{-1}$ ). Although ambergris is heterogenous (Baynes-Cope 1962), the metal contents of triplicate samples were quite reproducible herein (Figure S5). This metal profile suggests to us that the metal content of the flotsam reflects that of the liver, mantle, eyes and haemolymph of the squid in the sperm whale diet, rather than that of squid haemolymph alone (cf Lederer 1950).

Examination of a wider collection of samples (e.g. those studied by Rowland et al. 2019), will now allow this theory to be tested further.

### 3. Experimental

The experimental details are provided in the [Supplementary Material](#).

### 4. Conclusions

Analysis by multiple methods of a large piece of rare flotsam ambergris found in the Atlantic in 2019 produced data consistent with a post 1950s origin from a sperm whale. The metals profile was similar to that of the 'whole' body (e.g. liver, mantle, eyes and haemolymph) of the squid in the sperm whale diet, rather than that of squid haemolymph alone.

### Acknowledgments

We thank J. Quinn, University of Plymouth, for help with production of the graphical abstract (photos in graphical abstract: E.Berninsone/ARDITI and F. Alves).

### Disclosure statement

No potential conflict of interest was reported by the author(s).

### Funding

The Portuguese Foundation for Science and Technology (FCT) supported FA throughout the strategic projects UIDB/04292/2020 awarded to MARE and LA/P/0069/2020 granted to the Associate Laboratory ARNET.

### ORCID

Steven J. Rowland  <http://orcid.org/0000-0003-4980-0618>

Paul A. Sutton  <http://orcid.org/0000-0003-0568-5478>

C. Anthony Lewis  <http://orcid.org/0000-0002-0524-2215>

Timothy Knowles  <http://orcid.org/0000-0003-4871-5542>

Michael J. Wilde  <http://orcid.org/0000-0003-0726-890X>

Filipe Alves  <http://orcid.org/0000-0003-3752-2745>

Robert Clough  <http://orcid.org/0000-0003-3589-6671>

### References

- Baynes-Cope AD. 1962. Analyses of samples of ambergris. *Nature*. 193:978–979. doi:[10.1038/193978a0](https://doi.org/10.1038/193978a0).
- Brito CJ, Jordao VL, Pierce GJ. 2016. Ambergris as an overlooked historical marine resource: its biology and role as a global economic commodity. *J Mar Biol Assoc United Kingdom*. 96:585–596.
- Clarke R. 1954. A great haul of ambergris. *Nature*. 174:155–156. doi:[10.1038/174155a0](https://doi.org/10.1038/174155a0).

- Clarke R. 2006. The origin of ambergris. LAJAM. 5:7–21.
- Cornon R. 1955. Les odeurs ambrées. Industries de la Parfumerie. 10:291–295.
- Governo TF, Alessandro RT, Pragger MJ. 1977. Gas-liquid chromatographic-mass spectrometric detection and identification of ambergris. J AOAC Int. 60:160–164. doi:[10.1093/jaoac/60.1.160](https://doi.org/10.1093/jaoac/60.1.160).
- Lederer E. 1950. Odeurs et parfums des animaux. In: Zechmeister L, editor. Progress in the chemistry of organic natural products. Vol. 6. Vienna: springer-Verlag; p. 87–153.
- Macleod R, Sinding M-HS, Olsen MT, Collins MJ, Rowland SJ. 2020. DNA preserved in jetsam whale ambergris. Biol Lett. 16(2):20190819. doi:[10.1098/rsbl.2019.0819](https://doi.org/10.1098/rsbl.2019.0819).
- Mazzariol S, Di Guardo G, Petrella A, Marsili L, Fossi CM, Leonzio C, Zizzo N, Vizzini S, Gaspari S, Pavan G, et al. 2011. Sometimes sperm whales (*Physeter macrocephalus*) cannot find their way back to the high seas: a multidisciplinary study on a mass stranding. PLoS One. 6(5):e19417. doi:[10.1371/journal.pone.0019417](https://doi.org/10.1371/journal.pone.0019417).
- Mikhalev Y. 2014. Occurrence of sperm whale ambergris in mined flotilla "Slava" and "Soviet Ukraine. Ukrain Antarc J. 13:175–184. doi:[10.33275/1727-7485.13.2014.225](https://doi.org/10.33275/1727-7485.13.2014.225).
- Ohloff G, Schulte-Elte KH, Müller BL. 1977. Formation of ambergris odorants from ambrein under simulated natural conditions. Helv Chim Acta. 60:2763–2766.
- Rowland SJ, Payne D, Sutton PA. 2018a. Simulated environmental photo- and auto-oxidation of ambrein. <https://eartharxiv.org>.
- Rowland SJ, Sutton PA, Belt ST, Fitzsimmons-Thoss V, Scarlett AG. 2018. Further spectral and chromatographic studies of ambergris. Nat Prod Res. 32(21):2603–2609. doi:[10.1080/14786419.2018.1428599](https://doi.org/10.1080/14786419.2018.1428599).
- Rowland SJ, Sutton PA, Knowles TM. 2019. The age of ambergris. Nat Prod Res. 33:3134–3142.
- Rowland SJ, Sutton PA, Wolff GA. 2021. Biosynthesis of ambrein in ambergris: evidence from isotopic data and identification of possible intermediates. Nat Prod Res. 35(8):1235–1241. doi:[10.1080/14786419.2019.1644630](https://doi.org/10.1080/14786419.2019.1644630).
- Rowland SJ, Sutton PA. 2017. Chromatographic and spectral studies of jetsam and archived ambergris. Nat Prod Res. 31(15):1752–1757. doi:[10.1080/14786419.2017.1290618](https://doi.org/10.1080/14786419.2017.1290618).
- Rowland SJ, Wilde MJ, Sutton PA, Blackbird SJ, Wolff G. 2024. If speed is of the essence: rapid analysis of ambergris by APCI compact mass spectrometry. Nat Prod Res. 28:1–7. doi:[10.1080/14786419.2024.2321496](https://doi.org/10.1080/14786419.2024.2321496).
- Wilde MJ, Robson WJ, Sutton PA, Rowland SJ. 2020. Volatile and semi-volatile components of jetsam ambergris. Nat Prod Res. 34(21):3048–3053. doi:[10.1080/14786419.2019.1607855](https://doi.org/10.1080/14786419.2019.1607855).